

Datasheet: MCA2504

BATCH NUMBER 155368

Description:	MOUSE ANTI HUMAN CD44
Specificity:	CD44
Other names:	H-CAM, PGP-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	Bu52
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			1/100 - 1/200
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	<p>UniProt: P16070 Related reagents</p> <p>Entrez Gene: 960 CD44 Related reagents</p>
Synonyms	LHR, MDU2, MDU3, MIC4
RRID	AB_808430
Specificity	<p>Mouse anti Human CD44 antibody, clone Bu52 recognizes the human CD44 cell surface antigen, a ~100 kDa glycoprotein widely expressed on human leucocytes, white matter of the brain and by some epithelial cells of the intestine and of the breast. Several isoforms of CD44 exist, including the predominant CD44H isoform detected in many normal tissues. Mouse anti Human CD44 antibody, clone Bu52 recognizes an epitope that is common to all CD44 isoforms.</p> <p>CD44 is a receptor for hyaluronic acid (HA) and is involved in cell-cell interactions, cell adhesion and migration. CD44 also participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing. CD44 expression may be up-regulated upon some carcinomas, and it has been speculated that this may be related to metastatic potential.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Takahashi, K. <i>et al.</i> (1996) Keratan sulfate modification of CD44 modulates adhesion to hyaluronate. J Biol Chem. 271 (16): 9490-6. 2. Belitsos, P.C. <i>et al.</i> (1990) Homotypic cell aggregation induced by anti-CD44(Pgp-1) monoclonal antibodies and related to CD44(Pgp-1) expression. J Immunol. 144 (5): 1661-70. 3. Katz, E. <i>et al.</i> (2012) Targeting of Rac GTPases blocks the spread of intact human breast cancer. Oncotarget. 3 (6): 608-19. 4. Noda, S. <i>et al.</i> (2019) Effect of cell culture density on dental pulp-derived mesenchymal stem cells with reference to osteogenic differentiation. Sci Rep. 9 (1): 5430. 5. Hanke-Roos, M. <i>et al.</i> (2017) CD44 mediates the catch-bond activated rolling of HEPG2Iso epithelial cancer cells on hyaluronan. Cell Adh Migr. 11 (5-6): 476-87. 6. Muerza-Cascante, M.L. <i>et al.</i> (2017) Endosteal-like extracellular matrix expression on melt electrospun written scaffolds. Acta Biomater. 52: 145-58. 7. Thorne, R.F. <i>et al.</i> (2020) Evaluating nuclear translocation of surface receptors: recommendations arising from analysis of CD44. Histochem Cell Biol. 153 (2): 77-87.

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2504 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M367067:200529'

Printed on 18 Jan 2024