

Datasheet: MCA2516

Description:	MOUSE ANTI HUMAN CD87
Specificity:	CD87
Other names:	UPAR, UPA-R
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	VIM-5
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen	▪			1/100
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	THP-1 cells (Human acute monocytic leukaemia cells).
External Database Links	<p>UniProt: Q03405 Related reagents</p> <p>Entrez Gene: 5329 PLAUR Related reagents</p>
Synonyms	MO3, UPAR
RRID	AB_961472
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.
Specificity	<p>Mouse anti Human CD87 antibody, clone VIM-5 recognizes human CD87, also known as urokinase plasminogen activator receptor (uPAR), a 50-65 kDa single chain glycosylphosphatidylinositol (GPI) anchored glycoprotein, expressed by granulocytes, tissue macrophages, monocytes, endothelial cells, fibroblasts and mesenchymal and epithelial tumour cells.</p> <p>CD87 specifically binds and activates urokinase plasminogen activator (uPA), a serine protease which converts the inactive receptor-bound proenzyme plasminogen into plasmin, triggering the proteolysis cascade of fibrinolysis. The interaction between uPAR and uPA is also known to play a role in cancer cell invasion and metastasis.</p> <p>Clone VIM-5 is reported to block urokinase binding.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
Histology Positive Control Tissue	Human tonsil.
References	<ol style="list-style-type: none"> 1. Sillaber, C. <i>et al.</i> (1997) Molecular and functional characterization of the urokinase receptor on human mast cells. J Biol Chem. 272 (12): 7824-32. 2. Gadd, S.J. <i>et al.</i> (1990) M5, a phosphoinositol-linked human myelomonocytic activation-associated antigen. Clin Exp Immunol. 80 (2): 252-6.
Further Reading	1. Mondino, A. & Blasi, F. (2004) uPA and uPAR in fibrinolysis, immunity and pathology. Trends Immunol. 25 (8): 450-5.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	18 months from date of despatch.
------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@680 , DyLight@800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Recommended Useful Reagents

[HISTAR DETECTION SYSTEM \(STAR3000A\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------	------------------	-------------------------------------------------------------------------------------------------------------------------------------------------	---------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M331509:180912'

Printed on 09 Feb 2021