

Datasheet: MCA2516GA

Description:	MOUSE ANTI HUMAN CD87
Specificity:	CD87
Other names:	UPAR, UPA-R
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	VIM-5
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/200
Immunohistology - Frozen	▪			1/100
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	THP-1 cells (Human acute monocytic leukaemia cells).

**External Database
Links**

UniProt:

[Q03405](#) [Related reagents](#)

Entrez Gene:

[5329](#) PLAUR [Related reagents](#)

Synonyms

MO3, UPAR

RRID

AB_915235

Fusion Partners

Spleen cells from immunised Balb/c mice were fused with cells of the X63.Ag8.653 myeloma cell line.

Specificity

Mouse anti Human CD87 antibody, clone VIM-5 recognizes human CD87, also known as urokinase plasminogen activator receptor (uPAR), a 50-65 kDa single chain glycosylphosphatidylinositol (GPI) anchored glycoprotein, expressed by granulocytes, tissue macrophages, monocytes, endothelial cells, fibroblasts and mesenchymal and epithelial tumour cells.

CD87 specifically binds and activates urokinase plasminogen activator (uPA), a serine protease which converts the inactive receptor-bound proenzyme plasminogen into plasmin, triggering the proteolysis cascade of fibrinolysis. The interaction between uPAR and uPA is also known to play a role in cancer cell invasion and metastasis.

Clone VIM-5 is reported to block urokinase binding.

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

**Histology Positive
Control Tissue**

Human tonsil.

References

1. Sillaber, C. *et al.* (1997) Molecular and functional characterization of the urokinase receptor on human mast cells. [J Biol Chem. 272 \(12\): 7824-32.](#)
2. Gadd, S.J. *et al.* (1990) M5, a phosphoinositol-linked human myelomonocytic activation-associated antigen. [Clin Exp Immunol. 80 \(2\): 252-6.](#)

Further Reading

1. Mondino, A. & Blasi, F. (2004) uPA and uPAR in fibrinolysis, immunity and pathology. [Trends Immunol. 25 \(8\): 450-5.](#)

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@680 , DyLight@800 , FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

Recommended Useful Reagents

[HISTAR DETECTION SYSTEM \(STAR3000A\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M367094:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)