

Datasheet: MCA2518A647

Description:	MOUSE ANTI HUMAN CD172a:Alexa Fluor® 647
Specificity:	CD172a
Other names:	SIRP ALPHA
Format:	ALEXA FLUOR® 647
Product Type:	Monoclonal Antibody
Clone:	15-414
Isotype:	IgG2a
Quantity:	100 TESTS/1ml

Product Details

RRID AB_1055626

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Human

Product Form Purified IgG conjugated to Alexa Fluor® 647 - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Alexa Fluor®647	650	665

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide (NaN₃)
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.05mg/ml

Immunogen Monocyte-derived dendritic cells.

External Database Links **UniProt:**

Entrez Gene:

[140885](#) SIRPA [Related reagents](#)

Synonyms	BIT, MFR, MYD1, PTPNS1, SHPS1, SIRP
-----------------	-------------------------------------

Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the X63-Ag8.653 myeloma cell line.
------------------------	---

Specificity	<p>Mouse anti Human CD172a antibody, clone 15-414 recognizes human CD172a, also known as signal-regulatory protein alpha, a receptor-type transmembrane glycoprotein expressed on cells of myeloid origin, including granulocytes, dendritic cells (DCs), macrophages, mast cells and haematopoietic stem cells.</p> <p>CD172a acts as a substrate for several activated tyrosine kinases, including EGFR, PDGFR, src and insulin receptor and is involved in the negative regulation of receptor tyrosine kinase-coupled signaling pathways. Ligand binding of CD172a to integrin-associated protein CD47, results in tyrosine kinase phosphorylation of immunoreceptor tyrosine-based inhibitory motifs (ITIMs) within the cytoplasmic region of CD172a, mediating the recruitment and activation of the tyrosine phosphatases SHP-1 and SHP-2. These then act as regulators of cellular function, through dephosphorylation of specific substrates. Ligation of CD172a with CD47 has been demonstrated in several regulatory processes, including the inhibition of host cell phagocytosis by macrophages and the bi-directional activation of T cells and DCs.</p>
--------------------	---

Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
-----------------------	---

References	<ol style="list-style-type: none">1. Fujioka, Y. <i>et al.</i> (1996) A novel membrane glycoprotein, SHPS-1, that binds the SH2-domain-containing protein tyrosine phosphatase SHP-2 in response to mitogens and cell adhesion. Mol Cell Biol. 16 (12): 6887-99.2. Baba, T. <i>et al.</i> (2012) Novel Process of Intrathymic Tumor-Immune Tolerance through CCR2-Mediated Recruitment of Sirpα(+) Dendritic Cells: A Murine Model. PLoS One. 7: e41154.3. Szaraz, P.<i>et al.</i> (2016) <i>In Vitro</i> Differentiation of First Trimester Human Umbilical Cord Perivascular Cells into Contracting Cardiomyocyte-Like Cells Stem Cells International. 2016: 1-13.4. Hussen, J. <i>et al.</i> (2014) The chemokine CCL5 induces selective migration of bovine classical monocytes and drives their differentiation into LPS-hyporesponsive macrophages <i>in vitro</i>. Dev Comp Immunol. 47 (2): 169-77.
-------------------	--

Further Reading	1. van Beek, E.M. <i>et al.</i> (2005) Signal regulatory proteins in the immune system. J Immunol. 175 (12): 7781-7.
------------------------	--

Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>Storage in frost-free freezers is not recommended.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	--

Guarantee	18 months from date of despatch.
------------------	----------------------------------

Acknowledgements	This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in
-------------------------	--

research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or outlicensing@thermofisher.com

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA929A647\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M342790:190110'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)