

Datasheet: MCA2521

Description:	MOUSE ANTI HUMAN CD92
Specificity:	CD92
Other names:	CTL1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	VIM-15b
Isotype:	IgG2b
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	MV4-11 acute monocyte leukaemia cells.

**External Database
Links**

UniProt:

[Q8WWI5](#) [Related reagents](#)

Entrez Gene:

[23446](#) SLC44A1 [Related reagents](#)

Synonyms

CD92, CDW92, CTL1

RRID

AB_961480

Fusion Partners

Spleen cells from immunised Balb/c mice were fused with cells of the X63-Ag8.653 myeloma cell line.

Specificity

Mouse anti human CD92 antibody, clone VIM-15b recognizes the C-terminal variant of human CDw92 (isoform 3), also known as CTL1, a 70 kDa multi-pass membrane protein, expressed by monocytes, neutrophils, certain myeloid and T cell lines, and weakly by endothelial cells, fibroblasts and epithelial cells.

CDw92 is a member of the choline transporter-like protein family, so called due to their involvement in the efficient supply/transport of the natural amine choline, a vital cell nutrient required for the synthesis of cell membrane phospholipid components and the neurotransmitter acetylcholine.

Clone VIM-15b is reported to augment the LPS-induced production of IL-10 by monocyte-derived dendritic cells (Mo-DCs), and the reduced expression of CDw92 by Mo-DCs treated with ionomycin or calcium ionophore, can be reinduced in the presence of IL-10 ([Wille, S. et al.](#)). Removal of Sodium Azide is recommended prior to use in functional assays. Bio-Rad recommend the use of [EQU003](#) for this purpose.

Flow Cytometry

Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

**Histology Positive
Control Tissue**

Human tonsil.

References

1. Wille, S. *et al.* (2001) Characterization of CDw92 as a member of the choline transporter-like protein family regulated specifically on dendritic cells. [J Immunol. 167 \(10\): 5795-804.](#)
2. Majdic, O. *et al.* (1995) Leucocyte Typing V. Oxford University Press, Oxford pp.984-5.

Further Reading

1. O'regan, S. *et al.* (2000) An electric lobe suppressor for a yeast choline transport mutation belongs to a new family of transporter-like proteins. [Proc Natl Acad Sci U S A. 97 \(4\): 1835-40.](#)

Storage

Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

**Health And Safety
Information**

Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®680](#),
[DyLight®800](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL \(MCA691\)](#)

Recommended Useful Reagents

[HISTAR DETECTION SYSTEM \(STAR3000A\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M367108:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)