

Datasheet: MCA2537

| | |
|----------------------|-----------------------|
| Description: | MOUSE ANTI HUMAN CD16 |
| Specificity: | CD16 |
| Other names: | FcRIII |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | DJ130c |
| Isotype: | IgG1 |
| Quantity: | 0.2 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 1/50 - 1/200 |
| Immunohistology - Frozen | ▪ | | | |
| Immunohistology - Paraffin (1) | ▪ | | | |
| ELISA | | | ▪ | |
| Immunoprecipitation | | | ▪ | |
| Western Blotting | | | ▪ | |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species

Human

Species Cross Reactivity

Reacts with: Macaque

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG - liquid

| | |
|--|---|
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) |
| Carrier Free | Yes |
| Approx. Protein Concentrations | IgG concentration 1.0mg/ml |
| External Database Links | <p>UniProt:</p> <p>P08637 Related reagents</p> <p>O75015 Related reagents</p> <p>Entrez Gene:</p> <p>2214 FCGR3A Related reagents</p> <p>2215 FCGR3B Related reagents</p> |
| Synonyms | CD16A, CD16B, FCG3, FCGR3, IGFR3 |
| RRID | AB_877454 |
| Specificity | <p>Mouse anti Human CD16 antibody, clone DJ130c recognizes human CD16, also known as Low affinity immunoglobulin gamma Fc region receptor III-A or Fc-gamma RIIIa. CD16a is a 254 amino acid ~50-65 kDa single pass type 1 transmembrane glycoprotein bearing two Ig-like C2 type domains. CD16 exists as a transmembranous form (Fc gammaRIIIA, or CD16A) and a glycosyl phosphatidylinositol (GPI) anchored form, Fc gammaRIIIB, or CD16B (Scallon et al. 1989). CD16A is expressed by NK cells, some T cells, and macrophages, whereas CD16B is primarily expressed by granulocytes (Ravetch and Perussia 1989). In addition, CD16B exists as two allelic variants NA1 and NA2 . DJ130c recognizes all polymorphonuclear cells irrespective of their NA phenotype.</p> <p>Mouse anti Human CD16 antibody, clone DJ130c recognizes an epitope in the first membrane-distal domain of CD16, recognizes both CD16a and CD16b and has been demonstrated to cross-react with CD16 from rhesus macaques, <i>Macaca mulatta</i> (Xu et al. 2012)</p> |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. |
| Histology Positive Control Tissue | Human tonsil |
| References | 1. Schmidt, R.E. (1993) CD16 cluster workshop report. In Leucocyte Typing V: White cell differentiation antigens, Vol.1. Edited by Schlossman, S.F. <i>et al.</i> Oxford University Press. |

p805 – 806.

2. Kakko, T. *et al.* (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. [Diab Vasc Dis Res. 8: 221-8.](#)
3. Shantsila, E. *et al.* (2012) Fibrinolytic status in acute coronary syndromes: evidence of differences in relation to clinical features and pathophysiological pathways. [Thromb Haemost. 108: 32-40.](#)
4. Shantsila, E. *et al.* (2011) Immunophenotypic characterization of human monocyte subsets: possible implications for cardiovascular disease pathophysiology. [J Thromb Haemost. 9: 1056-66.](#)
5. Tapp, L.D. *et al.* (2012) The CD14⁺⁺CD16⁺ monocyte subset and monocyte-platelet interactions in patients with ST-elevation myocardial infarction. [J Thromb Haemost. 10: 1231-41.](#)
6. Ambarus, C.A. *et al.* (2012) Intimal lining layer macrophages but not synovial sublining macrophages display an IL-10 polarized-like phenotype in chronic synovitis. [Arthritis Res Ther. 14: R74.](#)
7. Ambarus, C.A. *et al.* (2012) Systematic validation of specific phenotypic markers for in vitro polarized human macrophages. [J Immunol Methods. 375: 196-206.](#)
8. Ambarus, C.A. *et al.* (2012) Soluble immune complexes shift the TLR-induced cytokine production of distinct polarized human macrophage subsets towards IL-10. [PLoS One. 7: e35994.](#)
9. Shantsila, E. *et al.* (2012) The effects of exercise and diurnal variation on monocyte subsets and monocyte-platelet aggregates. [Eur J Clin Invest. 42: 832-9.](#)
10. Chehadeh, W. *et al.* (2009) Antibody-mediated opsonization of red blood cells in parvovirus B19 infection. [Virology. 390: 56-63.](#)
11. Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. [Circ Heart Fail. 6: 127-35.](#)
12. Jaipersad, A.S. *et al.* (2014) Expression of monocyte subsets and angiogenic markers in relation to carotid plaque neovascularization in patients with pre-existing coronary artery disease and carotid stenosis. [Ann Med. 11: 1-9.](#)
13. Shantsila, E. *et al.* (2015) Free Light Chains in patients with acute coronary syndromes: Relationships to inflammation and renal function. [Int J Cardiol. 185: 322-7.](#)
14. Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. [Circ Heart Fail. 6 \(1\): 127-35.](#)
15. Romee R *et al.* (2013) NK cell CD16 surface expression and function is regulated by a disintegrin and metalloprotease-17 (ADAM17). [Blood. 121 \(18\): 3599-608.](#)
16. Sousa, S. *et al.* (2015) Human breast cancer cells educate macrophages toward the M2 activation status. [Breast Cancer Res. 17: 101.](#)
17. Shantsila, E. *et al.* (2019) Mon2 predicts poor outcome in ST-elevation myocardial infarction. [J Intern Med. 285 \(3\): 301-16.](#)
18. Brown, R.A. *et al.* (2018) Impact of Mon2 monocyte-platelet aggregates on human coronary artery disease. [Eur J Clin Invest. 48 \(5\): e12911.](#)
19. Nakajima-Kato, Y. *et al.* (2023) A novel monoclonal antibody with improved FcγR blocking ability demonstrated non-inferior efficacy compared to IVIG in cynomolgus monkey ITP model at considerably lower dose. [Clin Exp Immunol. 211 \(1\): 23-30.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at

-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

| | |
|------------------|---------------------------------|
| Guarantee | 12 months from date of despatch |
|------------------|---------------------------------|

| | |
|--------------------------------------|---|
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2537 10040 |
|--------------------------------------|---|

| | |
|-------------------|----------------------------|
| Regulatory | For research purposes only |
|-------------------|----------------------------|

Related Products

Recommended Secondary Antibodies

| | |
|---|---|
| Rabbit Anti Mouse IgG (STAR12...) | RPE |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | HRP |
| Goat Anti Mouse IgG (STAR76...) | RPE |
| Rabbit Anti Mouse IgG (STAR13...) | HRP |
| Goat Anti Mouse IgG (STAR70...) | FITC |
| Goat Anti Mouse IgG (H/L) (STAR117...) | Alk. Phos. , DyLight@488 , DyLight@550 , DyLight@650 , DyLight@680 , DyLight@800 , FITC , HRP |
| Rabbit Anti Mouse IgG (STAR9...) | FITC |
| Goat Anti Mouse IgG (STAR77...) | HRP |
| Goat Anti Mouse IgG (Fc) (STAR120...) | FITC , HRP |

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M383814:210513'

Printed on 18 Jan 2024