

## Datasheet: MCA2537A700

<b>Description:</b>	MOUSE ANTI HUMAN CD16:Alexa Fluor® 700
<b>Specificity:</b>	CD16
<b>Other names:</b>	FcRIII
<b>Format:</b>	ALEXA FLUOR® 700
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	DJ130c
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/1ml

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Macaque

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to Alexa Fluor® 700 - liquid

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
Alexa Fluor®700	702	723

#### Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

#### Buffer Solution

Phosphate buffered saline

<b>Preservative</b>	0.09% Sodium Azide
<b>Stabilisers</b>	1% Bovine Serum Albumin

<b>Approx. Protein Concentrations</b>	IgG concentration 0.05mg/ml
---------------------------------------	-----------------------------

<b>External Database Links</b>	<p><b>UniProt:</b></p> <p><a href="#">P08637</a>      <a href="#">Related reagents</a></p> <p><a href="#">O75015</a>      <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b></p> <p><a href="#">2214</a> FCGR3A    <a href="#">Related reagents</a></p> <p><a href="#">2215</a> FCGR3B    <a href="#">Related reagents</a></p>
--------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<b>Synonyms</b>	CD16A, CD16B, FCG3, FCGR3, IGFR3
-----------------	----------------------------------

<b>RRID</b>	AB_877457
-------------	-----------

**Specificity**      **Mouse anti Human CD16 antibody, clone DJ130c** recognizes human CD16, also known as Low affinity immunoglobulin gamma Fc region receptor III-A or Fc-gamma RIIIa. CD16a is a 254 amino acid ~50-65 kDa single pass type 1 transmembrane glycoprotein bearing two [Ig-like C2 type](#) domains. CD16 exists as a transmembranous form (Fc gammaRIIIA, or CD16A) and a glycosyl phosphatidylinositol (GPI) anchored form, Fc gammaRIIIB, or CD16B ([Scallon et al. 1989](#)). CD16A is expressed by NK cells, some T cells, and macrophages, whereas CD16B is primarily expressed by granulocytes ([Ravetch and Perussia 1989](#)). In addition, CD16B exists as two allelic variants NA1 and NA2 . DJ130c recognizes all polymorphonuclear cells irrespective of their NA phenotype.

Mouse anti Human CD16 antibody, clone DJ130c recognizes an epitope in the first membrane-distal domain of CD16, recognizes both CD16a and CD16b and has been demonstrated to cross-react with CD16 from rhesus macaques, *Macaca mulatta* ([Xu et al. 2012](#))

<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
-----------------------	---------------------------------------------------------------------------------------

<b>References</b>	<ol style="list-style-type: none"> <li>Schmidt, R.E. (1993) CD16 cluster workshop report. In Leucocyte Typing V: White cell differentiation antigens, Vol.1. Edited by Schlossman, S.F. <i>et al.</i> Oxford University Press. p805 – 806.</li> <li>Kakko, T. <i>et al.</i> (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. <a href="#">Diab Vasc Dis Res. 8: 221-8.</a></li> <li>Shantsila, E. <i>et al.</i> (2012) Fibrinolytic status in acute coronary syndromes: evidence of differences in relation to clinical features and pathophysiological pathways. <a href="#">Thromb Haemost. 108: 32-40.</a></li> <li>Shantsila, E. <i>et al.</i> (2011) Immunophenotypic characterization of human monocyte subsets: possible implications for cardiovascular disease pathophysiology. <a href="#">J Thromb</a></li> </ol>
-------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

[Haemost. 9: 1056-66.](#)

5. Tapp, L.D. *et al.* (2012) The CD14<sup>++</sup>CD16<sup>+</sup> monocyte subset and monocyte-platelet interactions in patients with ST-elevation myocardial infarction. [J Thromb Haemost. 10: 1231-41.](#)
6. Ambarus, C.A. *et al.* (2012) Intimal lining layer macrophages but not synovial sublining macrophages display an IL-10 polarized-like phenotype in chronic synovitis. [Arthritis Res Ther. 14: R74.](#)
7. Ambarus, C.A. *et al.* (2012) Systematic validation of specific phenotypic markers for in vitro polarized human macrophages. [J Immunol Methods. 375: 196-206.](#)
8. Ambarus, C.A. *et al.* (2012) Soluble immune complexes shift the TLR-induced cytokine production of distinct polarized human macrophage subsets towards IL-10. [PLoS One. 7: e35994.](#)
9. Shantsila, E. *et al.* (2012) The effects of exercise and diurnal variation on monocyte subsets and monocyte-platelet aggregates. [Eur J Clin Invest. 42: 832-9.](#)
10. Chehadeh, W. *et al.* (2009) Antibody-mediated opsonization of red blood cells in parvovirus B19 infection. [Virology. 390: 56-63.](#)
11. Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. [Circ Heart Fail. 6: 127-35.](#)
12. Jaipersad, A.S. *et al.* (2014) Expression of monocyte subsets and angiogenic markers in relation to carotid plaque neovascularization in patients with pre-existing coronary artery disease and carotid stenosis. [Ann Med. 11: 1-9.](#)
13. Shantsila, E. *et al.* (2015) Free Light Chains in patients with acute coronary syndromes: Relationships to inflammation and renal function. [Int J Cardiol. 185: 322-7.](#)
14. Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. [Circ Heart Fail. 6 \(1\): 127-35.](#)
15. Romee R *et al.* (2013) NK cell CD16 surface expression and function is regulated by a disintegrin and metalloprotease-17 (ADAM17). [Blood. 121 \(18\): 3599-608.](#)
16. Sousa, S. *et al.* (2015) Human breast cancer cells educate macrophages toward the M2 activation status. [Breast Cancer Res. 17: 101.](#)
17. Shantsila, E. *et al.* (2019) Mon2 predicts poor outcome in ST-elevation myocardial infarction. [J Intern Med. 285 \(3\): 301-16.](#)
18. Brown, R.A. *et al.* (2018) Impact of Mon2 monocyte-platelet aggregates on human coronary artery disease. [Eur J Clin Invest. 48 \(5\): e12911.](#)
19. Nakajima-Kato, Y. *et al.* (2023) A novel monoclonal antibody with improved FcγR blocking ability demonstrated non-inferior efficacy compared to IVIG in cynomolgus monkey ITP model at considerably lower dose. [Clin Exp Immunol. 211 \(1\): 23-30.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

**Guarantee**

12 months from date of despatch

---

**Acknowledgements**

This product is provided under an intellectual property licence from Life Technologies

Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com)

---

**Health And Safety Information**      Material Safety Datasheet documentation #10041 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA2537A700>  
10041

---

**Regulatory**                      For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 700 \(MCA928A700\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South**      Tel: +1 800 265 7376

**America**              Fax: +1 919 878 3751

                            Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

                            Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

                            Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M429751:240423'

**Printed on 23 Apr 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)