Datasheet: MCA2537BT BATCH NUMBER 151131

Description:	MOUSE ANTI HUMAN CD16:Biotin
Specificity:	CD16
Other names:	FcRIII
Format:	Biotin
Product Type:	Monoclonal Antibody
Clone:	DJ130c
lsotype:	lgG1
Quantity:	25 µg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	•			Neat - 1/10
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.				
Target Species	Human				
Species Cross Reactivity	Reacts with: Macaque N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.				
Product Form	Purified IgG conjugated	to Biotin - I	iquid		
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant				
Buffer Solution	Phosphate buffered salir	ıe			
Preservative	0.09% Sodium Azide (Na	aN ₃)			

Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		
External Database Links	UniProt: P08637 Related reagents O75015 Related reagents Entrez Gene: 2214 FCGR3A Related reagents 2215 FCGR3B Related reagents		
Synonyms	CD16A, CD16B, FCG3, FCGR3, IGFR3		
RRID	AB_1100855		
Specificity	 Mouse anti Human CD16 antibody, clone DJ130c recognizes human CD16, also known as Low affinity immunoglobulin gamma Fc region receptor III-A or Fc-gamma RIIIa. CD16a is a 254 amino acid ~50-65 kDa single pass type 1 transmembrane glycoprotein bearing two Ig-like C2 type domains. CD16 exists as a transmembranous form (Fc gammaRIIIA, or CD16A) and a glycosyl phosphatidylinositol (GPI) anchored form, Fc gammaRIIIB, or CD16B (Scallon <i>et al.</i> 1989). CD16A is expressed by NK cells, some T cells, and macrophages, whereas CD16B is primarily expressed by granulocytes (Ravetch and Perussia 1989). In addition, CD16B exists as two allelic variants NA1 and NA2. DJ130c recognizes all polymorphonuclear cells irrespective of their NA phenotype. Mouse anti Human CD16 antibody, clone DJ130c recognizes an epitope in the first membrane-distal domain of CD16, recognizes both CD16a and CD16b and has been demonstrated to cross-react with CD16 from rhesus macaques, <i>Macaca mulatta</i> (Xu <i>et al.</i> 2012) 		
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.		
References	 Schmidt, R.E. (1993) CD16 cluster workshop report. In Leucocyte Typing V: White cell differentiation antigens, Vol.1. Edited by Schlossman, S.F. <i>et al.</i> Oxford University Press. p805 – 806. Kakko, T. <i>et al.</i> (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. Diab Vasc Dis Res. 8: 221-8. Shantsila, E. <i>et al.</i> (2012) Fibrinolytic status in acute coronary syndromes: evidence of differences in relation to clinical features and pathophysiological pathways. Thromb Haemost. 108: 32-40. Shantsila, E. <i>et al.</i> (2011) Immunophenotypic characterization of human monocyte subsets: possible implications for cardiovascular disease pathophysiology. J Thromb Haemost. 9: 1056-66. 		

	5. Tapp, L.D. <i>et al.</i> (2012) The CD14++CD16+ monocyte subset and monocyte-platelet interactions in patients with ST-elevation myocardial infarction. <u>J Thromb Haemost. 10:</u>
	<u>1231-41.</u> 6. Ambarus, C.A. <i>et al.</i> (2012) Intimal lining layer macrophages but not synovial sublining
	macrophages display an IL-10 polarized-like phenotype in chronic synovitis. <u>Arthritis Res</u>
	<u>Ther. 14: R74.</u>
	7. Ambarus, C.A. et al. (2012) Systematic validation of specific phenotypic markers for in
	vitro polarized human macrophages. <u>J Immunol Methods. 375: 196-206.</u>
	8. Ambarus, C.A. et al. (2012) Soluble immune complexes shift the TLR-induced cytokine
	production of distinct polarized human macrophage subsets towards IL-10. <u>PLoS One. 7:</u>
	<u>e35994.</u> 9. Shantsila, E. <i>et al.</i> (2012) The effects of exercise and diurnal variation on monocyte
	subsets and monocyte-platelet aggregates. <u>Eur J Clin Invest. 42: 832-9.</u> 10. Chehadeh. W. <i>et al.</i> (2009) Antibody-mediated opsonization of red blood cells in
	parvovirus B19 infection. <u>Virology. 390: 56-63.</u>
	11. Wrigley, B.J. <i>et al.</i> (2013) Increased formation of monocyte-platelet aggregates in
	ischemic heart failure. <u>Circ Heart Fail. 6: 127-35.</u>
	12. Jaipersad, A.S. <i>et al.</i> (2014) Expression of monocyte subsets and angiogenic markers
	in relation to carotid plaque neovascularization in patients with pre-existing coronary artery
	disease and carotid stenosis. <u>Ann Med. 11: 1-9.</u> 13. Shantsila, E. <i>et al.</i> (2015) Free Light Chains in patients with acute coronary
	syndromes: Relationships to inflammation and renal function. Int J Cardiol. 185: 322-7.
	14. Wrigley, B.J. <i>et al.</i> (2013) Increased formation of monocyte-platelet aggregates in
	ischemic heart failure. <u>Circ Heart Fail. 6 (1): 127-35.</u>
	15. Romee R <i>et al.</i> (2013) NK cell CD16 surface expression and function is regulated by a
	disintegrin and metalloprotease-17 (ADAM17). <u>Blood. 121 (18): 3599-608.</u>
	16. Sousa, S. <i>et al.</i> (2015) Human breast cancer cells educate macrophages toward the
	M2 activation status. <u>Breast Cancer Res. 17: 101.</u>
	17. Shantsila, E. <i>et al.</i> (2019) Mon2 predicts poor outcome in ST-elevation myocardial
	infarction. <u>J Intern Med. 285 (3): 301-16.</u>
	18. Brown, R.A. <i>et al.</i> (2018) Impact of Mon2 monocyte-platelet aggregates on human
	coronary artery disease. <u>Eur J Clin Invest. 48 (5): e12911.</u>
Storage	Store at +4°C or at -20°C if preferred.
otorage	Storage in frost-free freezers is not recommended.
	This product should be stored undiluted. Avoid repeated freezing and thawing as this may
	denature the antibody. Should this product contain a precipitate we recommend
	microcentrifugation before use.
	microcentinugation belore use.
Guarantee	12 months from date of despatch
Health And Safety	Material Safety Datasheet documentation #10041 available at:
Information	https://www.bio-rad-antibodies.com/SDS/MCA2537BT
	10041
Populatory	
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M367135:200529'

Printed on 23 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint