

Datasheet: MCA2537PE

| Description: | MOUSE ANTI HUMAN CD16:RPE |
|---------------|---------------------------|
| Specificity: | CD16 |
| Other names: | FcRIII |
| Format: | RPE |
| Product Type: | Monoclonal Antibody |
| Clone: | DJ130c |
| Isotype: | IgG1 |
| Quantity: | 100 TESTS |
| | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | | | | Neat - 1/10 |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species | Human | | | | | |
|-----------------------------|---|--|--|--|--|--|
| Species Cross Reactivity | reactivity is derived | vity and working condit from testing within our I | ions may vary between species. Cross aboratories, peer-reviewed publications ors. Please refer to references indicated | | | |
| Product Form | Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized | | | | | |
| Reconstitution | Reconstitute with 1.0ml distilled water | | | | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) | | | |
| | RPE 488nm laser | 496 | 578 | | | |
| Preparation | Purified IgG prepare supernatant | ed by affinity chromatog | raphy on Protein A from tissue culture | | | |

| Buffer Solution | Phosphate buffered saline | | |
|-----------------------------|---|--|--|
| Preservative Stabilisers | 0.09% Sodium Azide (NaN₃)1% Bovine Serum Albumin5% Sucrose | | |
| External Database Links | UniProt: P08637 Related reagents O75015 Related reagents | | |
| | Entrez Gene: 2214 FCGR3A Related reagents 2215 FCGR3B Related reagents | | |
| Synonyms | CD16A, CD16B, FCG3, FCGR3, IGFR3 | | |
| RRID | AB_961449 | | |

Specificity

Mouse anti Human CD16 antibody, clone DJ130c recognizes human CD16, also known as Low affinity immunoglobulin gamma Fc region receptor III-A or Fc-gamma RIIIa. CD16a is a 254 amino acid ~50-65 kDa single pass type 1 transmembrane glycoprotein bearing two Ig-like C2 type domains. CD16 exists as a transmembranous form (Fc gammaRIIIA, or CD16A) and a glycosyl phosphatidylinositol (GPI) anchored form, Fc gammaRIIIB, or CD16B (Scallon et al. 1989). CD16A is expressed by NK cells, some T cells, and macrophages, whereas CD16B is primarily expressed by granulocytes (Ravetch and Perussia 1989). In addition, CD16B exists as two allelic variants NA1 and NA2.

Mouse anti Human CD16 antibody, clone DJ130c recognizes an epitope in the first membrane-distal domain of CD16, recognizes both CD16a and CD16b and has been demonstrated to cross-react with CD16 from rhesus macaques, *Macaca mulatta* (Xu et al. 2012)

Flow Cytometry

Use 10ul of the suggested working dilution to label 1x10⁶ cells in 100ul.

References

- 1. Schmidt, R.E. (1993) CD16 cluster workshop report. In Leucocyte Typing V: White cell differentiation antigens, Vol.1. Edited by Schlossman, S.F. *et al.* Oxford University Press. p805 806.
- 2. Kakko, T. *et al.* (2011) Inflammatory effects of blood leukocytes: association with vascular function in neuropeptide Y proline 7-genotyped type 2 diabetes patients. <u>Diab</u> Vasc Dis Res. 8: 221-8.
- 3. Shantsila, E. *et al.* (2012) Fibrinolytic status in acute coronary syndromes: evidence of differences in relation to clinical features and pathophysiological pathways. <u>Thromb</u> Haemost. 108: 32-40.
- 4. Shantsila, E. et al. (2011) Immunophenotypic characterization of human monocyte

subsets: possible implications for cardiovascular disease pathophysiology. <u>J Thromb</u> Haemost. 9: 1056-66.

- 5. Tapp, L.D. *et al.* (2012) The CD14++CD16+ monocyte subset and monocyte-platelet interactions in patients with ST-elevation myocardial infarction. <u>J Thromb Haemost. 10:</u> 1231-41
- 6. Ambarus, C.A. *et al.* (2012) Intimal lining layer macrophages but not synovial sublining macrophages display an IL-10 polarized-like phenotype in chronic synovitis. <u>Arthritis Res</u> Ther. 14: R74.
- 7. Ambarus, C.A. *et al.* (2012) Systematic validation of specific phenotypic markers for in vitro polarized human macrophages. <u>J Immunol Methods</u>. 375: 196-206.
- 8. Ambarus, C.A. *et al.* (2012) Soluble immune complexes shift the TLR-induced cytokine production of distinct polarized human macrophage subsets towards IL-10. <u>PLoS One. 7:</u> e35994.
- 9. Shantsila, E. *et al.* (2012) The effects of exercise and diurnal variation on monocyte subsets and monocyte-platelet aggregates. <u>Eur J Clin Invest. 42: 832-9.</u>
- 10. Chehadeh. W. *et al.* (2009) Antibody-mediated opsonization of red blood cells in parvovirus B19 infection. Virology. 390: 56-63.
- 11. Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. <u>Circ Heart Fail. 6: 127-35.</u>
- 12. Jaipersad, A.S. *et al.* (2014) Expression of monocyte subsets and angiogenic markers in relation to carotid plaque neovascularization in patients with pre-existing coronary artery disease and carotid stenosis. Ann Med. 11: 1-9.
- 13. Shantsila, E. *et al.* (2015) Free Light Chains in patients with acute coronary syndromes: Relationships to inflammation and renal function. <u>Int J Cardiol. 185: 322-7.</u>
- 14. Wrigley, B.J. *et al.* (2013) Increased formation of monocyte-platelet aggregates in ischemic heart failure. <u>Circ Heart Fail. 6 (1): 127-35.</u>
- 15. Romee R *et al.* (2013) NK cell CD16 surface expression and function is regulated by a disintegrin and metalloprotease-17 (ADAM17). <u>Blood. 121 (18): 3599-608</u>.
- 16. Sousa, S. *et al.* (2015) Human breast cancer cells educate macrophages toward the M2 activation status. Breast Cancer Res. 17: 101.
- 17. Shantsila, E. *et al.* (2019) Mon2 predicts poor outcome in ST-elevation myocardial infarction. J Intern Med. 285 (3): 301-16.
- 18. Brown, R.A. *et al.* (2018) Impact of Mon2 monocyte-platelet aggregates on human coronary artery disease. Eur J Clin Invest. 48 (5): e12911.

Storage

Store at +4°C.

DO NOT FREEZE.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody.

This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2537PE

20487

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA928PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375513:210104'

Printed on 12 Jun 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint