

## Datasheet: MCA2538A647

<b>Description:</b>	MOUSE ANTI HUMAN CD79a:Alexa Fluor® 647
<b>Specificity:</b>	CD79a
<b>Other names:</b>	MB-1
<b>Format:</b>	ALEXA FLUOR® 647
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	HM57
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/1ml

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/5 - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**(1)Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

<b>Target Species</b>	Human		
<b>Species Cross Reactivity</b>	Reacts with: Mouse, Rabbit, Horse, Pig, Monkey, Rat, Bovine, Guinea Pig, Fallow deer, American Bison, Red deer, Ferret, Goat <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.		
<b>Product Form</b>	Purified IgG conjugated to Alexa Fluor® 647 - liquid		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	Alexa Fluor®647	650	665
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture		

supernatant

---

<b>Buffer Solution</b>	Phosphate buffered saline
------------------------	---------------------------

---

<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> ) 1% Bovine Serum Albumin
---------------------------------	---

---

<b>Approx. Protein Concentrations</b>	IgG concentration 0.05mg/ml
---------------------------------------	-----------------------------

---

<b>Immunogen</b>	Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1
------------------	--

---

<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P11912</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">973</a> CD79A <a href="#">Related reagents</a>
--------------------------------	---

---

<b>Synonyms</b>	IGA, MB1
-----------------	----------

---

<b>RRID</b>	AB_905977
-------------	-----------

---

<b>Fusion Partners</b>	Spleen cells from immunised Balb/c mice were fused with cells of the Sp2/0 myeloma cell line
------------------------	--

---

<b>Specificity</b>	<p><b>Mouse anti Human CD79a antibody, clone HM57</b> recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45 kDa protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells.</p> <p>The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signaling following antigen-binding to surface immunoglobulin.</p>
--------------------	--

---

<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
-----------------------	---

---

<b>References</b>	<ol style="list-style-type: none"><li>1. Mason, D.Y. <i>et al.</i> (1991) The IgM-associated protein mb-1 as a marker of normal and neoplastic B cells. <a href="#">J Immunol. 147 (11): 2474-82.</a></li><li>2. Jones, M. <i>et al.</i> (1993) Detection of T and B cells in many animal species using cross-reactive anti-peptide antibodies. <a href="#">J Immunol. 150 (12): 5429-35.</a></li><li>3. Christgau, M. <i>et al.</i> (1998) Characterization of immunocompetent cells in the diseased canine periodontium. <a href="#">J Histochem Cytochem. 46: 1443-54.</a></li><li>4. Spaas, J.H. <i>et al.</i> (2013) Culture and characterisation of equine peripheral blood mesenchymal stromal cells. <a href="#">Vet J. 195 (1): 107-13.</a></li><li>5. Nelson, D.D. <i>et al.</i> (2010) CD8(+)/perforin(+)/WC1(-) gammadelta T cells, not CD8(+) alphabeta T cells, infiltrate vasculitis lesions of American bison (<i>Bison bison</i>) with</li></ol>
-------------------	---

- experimental sheep-associated malignant catarrhal fever. [Vet Immunol Immunopathol. 136: 284-91.](#)
6. De Schauwer, C. *et al.* (2012) In search for cross-reactivity to immunophenotype equine mesenchymal stromal cells by multicolor flow cytometry. [Cytometry A. 81 \(4\): 312-23.](#)
7. Long, H. *et al.* (2016) Polyostotic Lymphoma in a Ferret (*Mustela putorius furo*). [J Comp Pathol. 154 \(4\): 341-4.](#)
8. Schinköthe J *et al.* (2016) Characterization of tuberculous granulomas in different stages of progression and associated tertiary lymphoid tissue in goats experimentally infected with *Mycobacterium avium* subsp. *hominissuis*. [Comp Immunol Microbiol Infect Dis. 47: 41-51.](#)
9. Bozkurt, Y.A., *et al.* (2014) Histological and immunohistological studies of the structure of lymph nodes in Kilis goats. [Biotech Histochem. 89\(6\):440-5.](#)
10. Froment R & Bédard C (2016) Marked hyperphosphatasemia associated with an acute leukemia in a Great Dane. [Vet Clin Pathol. Aug 18. \[Epub ahead of print\]](#)
11. Aresu, L. *et al.* (2015) Canine indolent and aggressive lymphoma: clinical spectrum with histologic correlation. [Vet Comp Oncol. 13 \(4\): 348-62.](#)
12. Poggi, A. *et al.* (2015) Flow cytometric evaluation of ki67 for the determination of malignancy grade in canine lymphoma. [Vet Comp Oncol. 13 \(4\): 475-80.](#)
13. Gelain ME *et al.* (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. [Vet Immunol Immunopathol. 159 \(1-2\): 91-6.](#)
14. Paebst, F. *et al.* (2014) Comparative immunophenotyping of equine multipotent mesenchymal stromal cells: an approach toward a standardized definition. [Cytometry A. 85 \(8\): 678-87.](#)
15. De Schauwer, C. *et al.* (2014) Characterization and profiling of immunomodulatory genes of equine mesenchymal stromal cells from non-invasive sources. [Stem Cell Res Ther. 5 \(1\): 6.](#)
16. Claessen, C. *et al.* (2015) Equid herpesvirus 1 (EHV1) infection of equine mesenchymal stem cells induces a pUL56-dependent downregulation of select cell surface markers. [Vet Microbiol. 176 \(1-2\): 32-9.](#)
17. Novacco, M. *et al.* (2015) Prognostic factors in canine acute leukaemias: a retrospective study. [Vet Comp Oncol. Jan 26. \[Epub ahead of print\]](#)
18. Hillmann, A. *et al.* (2016) Comparative Characterization of Human and Equine Mesenchymal Stromal Cells: A Basis for Translational Studies in the Equine Model. [Cell Transplant. 25 \(1\): 109-24.](#)
19. Moore, P.F. *et al.* (2013) Canine inflamed nonepitheliotropic cutaneous T-cell lymphoma: a diagnostic conundrum. [Vet Dermatol. 24 \(1\): 204-11.e44-5.](#)
20. Nagata, K. *et al.* (2017) Epstein-Barr Virus Lytic Reactivation Activates B Cells Polyclonally and Induces Activation-Induced Cytidine Deaminase Expression: A Mechanism Underlying Autoimmunity and Its Contribution to Graves' Disease. [Viral Immunol. Mar 23. \[Epub ahead of print\]](#)
21. Wessels, M. *et al.* (2017) Systemic necrotizing polyarteritis in three weaned lambs from one flock. [J Vet Diagn Invest. May 1:1040638717709856. \[Epub ahead of print\]](#)
22. Uitterdijk, A. *et al.* (2017) Time course of VCAM-1 expression in reperfused myocardial infarction in swine and its relation to retention of intracoronary administered bone marrow-derived mononuclear cells. [PLoS One. 12 \(6\): e0178779.](#)
23. Forner, R. *et al.* (2021) Distribution difference of colostrum-derived B and T cells subsets in gilts and sows. [PLoS One. 16 \(5\): e0249366.](#)

24. Murphy, E.G. *et al.* (2019) First detection of Hepatitis E virus (Orthohepevirus C) in wild brown rats (*Rattus norvegicus*.) from Great Britain. [Zoonoses Public Health. 66 \(6\): 686-94.](#)
25. Skovdal, S.M. *et al.* (2019) Inhaled nebulized glatiramer acetate against Gram-negative bacteria is not associated with adverse pulmonary reactions in healthy, young adult female pigs. [PLoS One. 14 \(10\): e0223647.](#)
26. Collins, J.J.P. *et al.* (2018) Impaired Angiogenic Supportive Capacity and Altered Gene Expression Profile of Resident CD146<sup>+</sup> Mesenchymal Stromal Cells Isolated from Hyperoxia-Injured Neonatal Rat Lungs. [Stem Cells Dev. 27 \(16\): 1109-24.](#)
27. Matsuyama, S. *et al.* (2021) Properties of macrophages and lymphocytes appearing in rat renal fibrosis followed by repeated injection of cisplatin. [J Vet Med Sci. 83 \(9\): 1435-42.](#)
28. Mu&tilde;oz-Silvestre, A. *et al.* (2020) Pathogenesis of Intradermal Staphylococcal Infections: Rabbit Experimental Approach to Natural *Staphylococcus aureus* Skin Infections. [Am J Pathol. 190 \(6\): 1188-210.](#)
29. Carroll, C.S.E. *et al.* (2021) Simple and effective bacterial-based intratumoral cancer immunotherapy. [J Immunother Cancer. 9 \(9\): e002688.\[Epub ahead of print\].](#)
30. do Prado Duzanski, A. *et al.* (2022) Cell-mediated immunity and expression of MHC class I and class II molecules in dogs naturally infected by canine transmissible venereal tumor: Is there complete spontaneous regression outside the experimental CTVT? [Res Vet Sci. 145: 193-204.](#)
31. Maciag, S.S. *et al.* (2022) The influence of source of porcine colostrum in development of early immune ontogeny in the piglet [Res Sq. Mar 24 \[Epub ahead of print\].](#)

---

**Further Reading** 1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

---

**Guarantee** 12 months from date of despatch

---

**Acknowledgements** This product is provided under an intellectual property licence from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchase product solely in research, excluding contract research or any fee for service research, and the buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad CA 92008 USA or [outlicensing@thermofisher.com](mailto:outlicensing@thermofisher.com)

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at:  
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:Alexa Fluor® 647 \(MCA928A647\)](#)

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M385453:210513'

**Printed on 11 Apr 2022**

---

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)