

Datasheet: MCA2539GA

Description:	RAT ANTI MOUSE PLVAP
Specificity:	PLVAP
Other names:	MECA-32, PLASMALEMMA VESICLE ASSOCIATED PROTEIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	MECA-32
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			1/25 - 1/100
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
External Database Links	<p>UniProt: Q91VC4 Related reagents</p> <p>Entrez Gene: 84094 Plvap Related reagents</p>
Synonyms	Pv1
RRID	AB_931734
Fusion Partners	Spleen cells from an immunized rat were fused with cells of the Sp2/0 myeloma cell line.
Specificity	<p>Rat anti Mouse PLVAP antibody, clone MECA-32 recognizes murine plasmalemma vesicle-associated protein (PLVAP), a ~60 kDa protein also known as MECA-32 and PV-1. PLVAP is a transmembrane homodimer associated with endothelial cell caveolae and fenestrae. In adult and embryonic mice, PLVAP is primarily expressed on endothelial tissues but shows restricted distribution in skeletal, cardiac and brain tissues. In embryonic mice, PLVAP expression, on the vasculature associated with blood brain barrier has been reported to be developmentally regulated.</p> <p>In humans, PLVAP is selectively up-regulated in a variety of high-grade brain tumors, including glioblastoma and metastatic carcinoma, as well as other cerebral disorders associated with blood-brain barrier disruption, such as acute ischemia. PLVAP represents a novel marker of brain tumour angiogenesis and integrity of the blood-brain barrier and is a potential therapeutic target.</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	<ol style="list-style-type: none"> Hallmann, R. <i>et al.</i> (1995) Novel mouse endothelial cell surface marker is suppressed during differentiation of the blood brain barrier. Dev Dyn. 202 (4): 325-32. Orford M <i>et al.</i> (2009) Generation of an ABCG2(GFPn-puro) transgenic line--a tool to study ABCG2 expression in mice. Biochem Biophys Res Commun. 384 (2): 199-203. Daneman, R. <i>et al.</i> (2010) Pericytes are required for blood-brain barrier integrity during embryogenesis. Nature. 468 (7323): 562-6. Okizaki, S. <i>et al.</i> (2016) Vascular Endothelial Growth Factor Receptor Type 1 Signaling Prevents Delayed Wound Healing in Diabetes by Attenuating the Production of IL-1β by Recruited Macrophages. Am J Pathol. 186 (6): 1481-98.
Further Reading	1. Carson-Walter, E.B. <i>et al.</i> (2005) Plasmalemmal vesicle associated protein-1 is a novel marker implicated in brain tumor angiogenesis. Clin Cancer Res. 11 (21): 7643-50.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2539GA 10040
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M383940:210513'

Printed on 18 Jan 2024