

Datasheet: MCA2558PE

BATCH NUMBER 155267

Description:	MOUSE ANTI DOG CD107b:RPE
Specificity:	CD107b
Other names:	LAMP-2
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	AC17
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.

Target Species

Dog

Species Cross Reactivity

Reacts with: Mink, Human

Does not react with: Mouse, Rat

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

Reconstitution

Reconstitute with 1ml distilled water

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)

Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin 5% Sucrose
Immunogen	MDCK (Madin-Darby Canine Kidney) cells
RRID	AB_10897185
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the NS1 myeloma cell line
Specificity	<p>Mouse anti Dog CD107b antibody, clone AC17 recognizes canine CD107b, otherwise known as lysosome-associated membrane protein 2 or LAMP-2. Immunofluorescence staining of MDCK cells with mouse anti dog CD107b, clone AC17 demonstrates staining patterns consistent with localization to lysosomes. This is supported by coincident staining of an exogenous lysosomal glycoprotein, avian LEP100 transfected into MDCK cells and detected using the anti LEP100 antibody clone CV24 (Nabi et al.1991).</p> <p>Mouse anti Dog CD107b antibody, clone AC17 immunoprecipitates a protein of ~95 kDa in MDCK cells which, following Endo F digestion to remove N-linked oligosaccharides, yields a core protein product of 40 kDa, indicating the heavily glycosylated nature of CD107b. The molecular weight of canine CD107b is typical of many lysosome-associated membrane proteins. While most (97%) CD107b resides in the lysosomal environment in adherent MDCK cells <i>in vitro</i>, a small percentage is associated with the cell membrane (Nabi et al.1991).</p> <p>CD107b has been shown to share high N-terminal amino acid sequence homology with human, mouse and rat CD107b (Nabi et al.1993).</p> <p>Transfection of a mink type II lung epithelial cell line with beta1-6-N-acetylglucosaminyl transferase V demonstrates the formation of large lysosomal vacuoles, termed multilamellar bodies (MLBs), having a very distinct phenotype with expression of CD107b, as indicated by immunofluorescent staining with clone AC17. These MLBs require lysosomal degradation via an autophagic pathway for their formation and may have implications for lysosomal storage diseases (Hariri et al.2000). CD107b is involved in the lysosomal uptake of cytosolic proteins and the endocytic pathway.</p> <p>Mouse anti Dog CD107b antibody, clone AC17 is suitable for use in electron microscopy (Nabi et al.1991).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. Nabi, I.R. *et al.* (1991) An endogenous MDCK lysosomal membrane glycoprotein is targeted basolaterally before delivery to lysosomes. [J Cell Biol. 115 \(6\): 1573-84.](#)
2. Nabi, I.R. & Rodriguez-Boulan, E. (1993) Increased LAMP-2 polylectosamine glycosylation is associated with its slower Golgi transit during establishment of a polarized MDCK epithelial monolayer. [Mol Biol Cell. 4 \(6\): 627-35.](#)
3. Jou, T.S. *et al.* (2000) Selective alterations in biosynthetic and endocytic protein traffic in Madin-Darby canine kidney epithelial cells expressing mutants of the small GTPase Rac1. [Mol Biol Cell. 11 \(1\): 287-304.](#)
4. Ihrke, G. *et al.* (2001) Competing sorting signals guide endolyn along a novel route to lysosomes in MDCK cells. [EMBO J. 20 \(22\): 6256-64.](#)
5. Pluhar, G.E. *et al.* (2010) Anti-tumor immune response correlates with neurological symptoms in a dog with spontaneous astrocytoma treated by gene and vaccine therapy. [Vaccine 28 \(19\): 3371-8.](#)
6. Cliffe, S.T. *et al.* (2009) SLC29A3 gene is mutated in pigmented hypertrichosis with insulin-dependent diabetes mellitus syndrome and interacts with the insulin signaling pathway. [Hum Mol Genet. 18: 2257-65.](#)
7. Bai, Y. *et al.* (2011) Intracellular neutralization of viral infection in polarized epithelial cells by neonatal Fc receptor (FcRn)-mediated IgG transport. [Proc Natl Acad Sci U S A. 108 \(45\): 18406-11.](#)
8. Nagahama, M. *et al.* (2011) Cellular vacuolation induced by *Clostridium perfringens* epsilon-toxin. [FEBS J. 278: 3395-407.](#)
9. Nagahama, M. *et al.* (2012) Intracellular trafficking of *Clostridium perfringens* iota-toxin b. [Infect Immun. 80: 3410-6.](#)
10. Hariri, M. *et al.* (2000) Biogenesis of multilamellar bodies via autophagy. [Mol Biol Cell. 11: 255-68.](#)

Further Reading

1. Fukuda, M. (1991) Lysosomal membrane glycoproteins. Structure, biosynthesis, and intracellular trafficking. [J Biol Chem. 266 \(32\): 21327-30.](#)

Storage

Prior to reconstitution store at +4°C.
After reconstitution store at +4°C.
DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #20487 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2558PE>
20487

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:RPE \(MCA928PE\)](#)

North & South Tel: +1 800 265 7376

Worldwide Tel: +44 (0)1865 852 700

Europe Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M375516:210104'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)