

# Datasheet: MCA2576 BATCH NUMBER 150982

Description:	MOUSE ANTI ASPERGILLUS SPP		
Specificity:	ASPERGILLUS SPP		
Format:	Purified		
<b>Product Type:</b>	Monoclonal Antibody		
Clone:	WF-AF-1		
Isotype:	IgM		
Quantity:	0.25 mg		

## **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen				
Immunohistology - Paraffin (1)	•			1/300
Immunoprecipitation				
Western Blotting	-			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires protein digestion pre-treatment of paraffin sections e.g. See <u>Jensen et al.</u> (2000) for details.

Target Species	Fungal	
Product Form	Purified IgM - liquid	
Preparation	Purified IgM prepared by ammonium sulfate precipitation from	n tissue culture supernatant
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )	

Approx. Protein Concentrations	IgM concentration 1.0 mg/ml
Immunogen	Wall fraction (WF) of Aspergillus fumigatus
RRID	AB_1100465
Fusion Partners	Spleen cells from immunised Balb/c ABom mice were fused with cells of the X63-Ag8.653 myeloma cell line.
Specificity	<b>Mouse anti</b> <i>Aspergillus spp.</i> , <b>clone WF-AF-1</b> , is raised against the wall fraction (WF) of <i>Aspergillus fumigatus</i> . This antibody specifically recognizes members of the <i>Aspergillus spp</i> . including <i>A. flavus</i> and <i>A. niger</i> , reacting strongly with walls and septae, and to a lesser extent within the cytoplasm of hyphae.
	A. fumigatus, a thermophilic, opportunistic and angio-invasive filamentous fungus, is the main causative agent of systemic bovine aspergillosis, a worldwide and often fatal respiratory disease of cattle. Clone WF-AF-1 has been successfully used in immunohistochemistry for the specific and consistent in situ diagnosis of bovine systemic aspergillosis, attributed to its binding to the major cell wall component, galactomannan. Clone WF-AF-1 has also been used for the identification of aspergillosis in human tissue sections.
	Mouse anti Aspergillus spp., clone WF-AF-1, does not bind to water-soluble somatic antigens (WSSA) of Aspergillus spp., but may react with galactomannans of members of the genus <i>Penicillium</i> .
Histology Positive Control Tissue	Aspergillus infected placenta.
Western Blotting	Mouse anti <i>Aspergillus</i> spp. antibody, clone WF-AF-1 detects a band of approximately 106kDa of <i>Aspergillus fumigatus</i> wall fraction (WF).
References	<ol> <li>Jensen, H.E. <i>et al.</i> (1996) Development of murine monoclonal antibodies for the immunohistochemical diagnosis of systemic bovine aspergillosis. J Vet Diagn Invest. 8 (1): 68-75.</li> <li>Jensen, H.E. <i>et al.</i> (1996) Diagnosis of systemic mycoses by specific immunohistochemical tests. APMIS. 104 (4): 241-58.</li> <li>Jensen, H.E. <i>et al.</i> (1997) The use of immunohistochemistry to improve sensitivity and specificity in the diagnosis of systemic mycoses in patients with haematological malignancies. J Pathol. 181 (1): 100-5.</li> <li>Delaney, M.A. <i>et al.</i> (2013) Occlusive fungal tracheitis in 4 captive bottlenose dolphins (<i>Tursiops truncatus</i>). Vet Pathol. 50 (1): 172-6.</li> <li>Goodpaster, T. &amp; Randolph-Habecker, J. (2014) A flexible mouse-on-mouse immunohistochemical staining technique adaptable to biotin-free reagents, immunofluorescence, and multiple antibody staining. J Histochem Cytochem. 62 (3): 197-204.</li> </ol>

6. Galiza Glauco J.N. et al. (2014) Usage of three immunohistochemical methods in the

detection of aspergillosis and zygomycosis in animals Pesquisa Veterinária Brasileira. 34

- 7. Murase, H. et al. (2015) A clinical case of equine fungal placentitis with reference to hormone profiles and ultrasonography. <u>J Equine Sci. 26 (4): 129-33.</u>
- 8. Dagleish, M.P. et al. (2010) Immunohistochemical diagnosis of infectious diseases of sheep. Small Ruminant Research. 92 (1-3): 19-35.
- 9. Suzuta F et al. (2015) Variations in the morphology of Rhizomucor pusillus in granulomatous lesions of a Magellanic penguin (Spheniscus magellanicus). J Vet Med Sci. 77 (8): 1029-31.
- 10. Jin, J-H. et al. (2015) Real-time selective monitoring of allergenic Aspergillus molds using pentameric antibody-immobilized single-walled carbon nanotube-field effect transistors RSC Adv. 5 (20): 15728-15735.
- 11. Ogasawara, F. et al. (2016) Concurrent Fowlpox and Candidiasis Diseases in Backyard Chickens with Unusual Pox Lesions in the Bursa of Fabricius. Avian Dis. 60 (3): <del>705-8.</del>

#### **Storage**

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2576">https://www.bio-rad-antibodies.com/SDS/MCA2576</a> 10040
Regulatory	For research purposes only

## Related Products

## **Recommended Secondary Antibodies**

Goat Anti Mouse IgM (STAR138...) Alk. Phos. Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

America

North & South Tel: +1 800 265 7376 Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody sales us@bio-rad.com

Email: antibody sales uk@bio-rad.com

Email: antibody sales de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M367194-200529'

### Printed on 19 Jan 2024