

Datasheet: MCA2578F

Description:	RAT ANTI MOUSE TREM-1:FITC
Specificity:	TREM-1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	L5-B8.2A12.3A12
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

RRID AB_2256512

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Mouse

Product Form Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide (NaN₃)
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1mg/ml

Immunogen Mouse TREM-1 Fc fusion protein.

External Database Links

UniProt:
[Q9JKE2](https://www.uniprot.org/entry/Q9JKE2) [Related reagents](#)

Entrez Gene:[58217](#) Trem1 [Related reagents](#)

Specificity **Rat anti Mouse Trem-1 antibody, clone L5-B8.2A12.3A12** recognizes mouse triggering receptor expressed on myeloid cells 1 (TREM-1), a 30kD member of the immunoglobulin superfamily expressed on neutrophils and a subset of monocytes. TREM-1 is involved in the microbial inflammatory response. It acts in conjunction with the adaptor protein DAP12 to activate neutrophils and monocytes, leading to the secretion of pro-inflammatory mediators and enhancement of the inflammatory response. Microbial infection also induces the release of soluble form of TREM-1 from monocytes, which is reported to act as an inhibitor of TREM-1.

TREM-1 mediates the septic shock pathway, and blocking TREM-1 has been reported to reduce inflammation and increase survival in certain models of bacterial infection.

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References 1. Wang, F. *et al.* (2012) Blocking TREM-1 signaling prolongs survival of mice with *Pseudomonas aeruginosa* induced sepsis. [Cell Immunol. 272 \(2\): 251-8.](#)

Storage Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL:FITC \(MCA1212F\)](#)

Recommended Useful Reagents

[MOUSE SEROBLOCK FcR \(BUF041A\)](#)

[MOUSE SEROBLOCK FcR \(BUF041B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M343681:190111'

Printed on 20 May 2019