

Datasheet: MCA2578PE

### **BATCH NUMBER 162191**

Description:	RAT ANTI MOUSE TREM-1:RPE
Specificity:	TREM-1
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	L5-B8.2A12.3A12
Isotype:	lgG2a
Quantity:	100 TESTS

## **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse				
Product Form	Purified IgG conjuga	ated to R. Phycoerythrin	ı (RPE) - Iyophilized		
Reconstitution	Reconstitute with 1.0	0ml distilled water			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	RPE 488nm laser	496	578		
Preparation	Purified IgG prepared by affinity chromatography on Protein G supernatant				
Buffer Solution	Phosphate buffered	saline			
Preservative	0.09% Sodium Azido	e			
Stabilisers	1% Bovine Serur	n Albumin			
	5% Sucrose				

Immunogen	Mouse TREM-1 Fc fusion protein.
External Database Links	UniProt:  Q9JKE2 Related reagents  Entrez Gene:  58217 Trem1 Related reagents
RRID	AB_10892677
Specificity	Rat anti Mouse Trem-1 antibody, clone L5-B8.2A12.3A12 recognizes mouse triggering receptor expressed on myeloid cells 1 (TREM-1), a 30kD member of the immunoglobulin superfamily expressed on neutrophils and a subset of monocytes. TREM-1 is involved in the microbial inflammatory response. It acts in conjunction with the adaptor protein DAP12 to activate neutrophils and monocytes, leading to the secretion of pro-inflammatory mediators and enhancement of the inflammatory response. Microbial infection also induces the release of soluble form of TREM-1 from monocytes, which is reported to act as an inhibitor of TREM-1.
	TREM-1 mediates the septic shock pathway, and blocking TREM-1 has been reported to reduce inflammation and increase survival in certain models of bacterial infection.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
References	1. Wang, F. et al. (2012) Blocking TREM-1 signaling prolongs survival of mice with Pseudomonas aeruginosa induced sepsis. Cell Immunol. 272 (2): 251-8.
Storage	Store at +4°C. DO NOT FREEZE.  This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2578PE">https://www.bio-rad-antibodies.com/SDS/MCA2578PE</a> 20487
Regulatory	For research purposes only

# **Related Products**

**Recommended Negative Controls** 

RAT IgG2a NEGATIVE CONTROL:RPE (MCA1212PE)

**Recommended Useful Reagents** 

MOUSE SEROBLOCK FcR (BUF041A)

### MOUSE SEROBLOCK FcR (BUF041B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375518:210104'

### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint