

# Datasheet: MCA2584F BATCH NUMBER 166774

Description:	MOUSE ANTI HUMAN CD172g:FITC			
Specificity:	CD172g			
Other names:	SIRP GAMMA, SIRPGAMMA			
Format:	FITC			
<b>Product Type:</b>	Monoclonal Antibody			
Clone:	OX119			
Isotype:	lgG1			
Quantity:	0.1 mg			

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

# **Immunogen** Purified recombinant SIRPgamma.CD4. **External Database UniProt:** Links **Q9P1W8** Related reagents **Entrez Gene:** 55423 SIRPG Related reagents **Synonyms** SIRPB2 **RRID** AB 961530 **Fusion Partners** Spleen cells from immunised BALB/c mice were fused with cells of the NS-1 myeloma cell line. **Specificity** Mouse anti Human CD172g antibody, clone OX119 recognizes human SIRP gamma, a 55 kDa signal regulatory protein which is a member of the immunoglobulin gene superfamily. SIRP gamma, also known as CD172g, is expressed on most T lymphocytes and a subset of B-cells but is absent on myeloid cells. CD172g has a truncated cytoplasmic tail, which is similar to SIRP beta, but unlike SIRP beta CD172g does not require DAP12 for expression at the cell surface. CD172q, like SIRP alpha, binds to CD47 inducing apoptosis. It is also involved in the negative regulation of receptor tyrosine kinase-coupled signalling. **Flow Cytometry** Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul. References 1. Brooke, G. et al. (2004) Human lymphocytes interact directly with CD47 through a novel member of the signal regulatory protein (SIRP) family. J Immunol.173:2562-70. **Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Guarantee 12 months from date of despatch **Health And Safety** Material Safety Datasheet documentation #10041 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA2584F 10041 Regulatory For research purposes only

#### Related Products

Concentrations

### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

# **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M422726:230928'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint