

Datasheet: MCA2588GA

Description:	MOUSE ANTI HUMAN CD61
Specificity:	CD61
Other names:	INTEGRIN BETA 3 CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	Y2/51
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			
ELISA			▪	
Immunoprecipitation			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin prior to staining.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)

Stabilisers

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0mg/ml

Immunogen PHA stimulated peripheral blood cells.

External Database Links

UniProt:

[P05106](#) [Related reagents](#)

Entrez Gene:

[3690](#) ITGB3 [Related reagents](#)

Synonyms GP3A

RRID AB_1055675

Fusion Partners Spleen cells from immunised Balb/c mice were fused with cells of the NS1 myeloma cell line.

Specificity **Mouse anti Human CD61 antibody, clone Y2/51** recognizes CD61, a ~105 kDa glycoprotein, also known as integrin beta 3 chain. This molecule associates with either the alpha IIb integrin (CD41) or the alpha V integrin (CD51) at the cell surface. CD61 is expressed on platelets and megakaryocytes in association with CD41, and on endothelial cells, monocytes, platelets and osteoclasts in association with CD51.

CD61 is a receptor for fibrinogen, fibronectin, vWF, vitronectin and thrombospondin.

Flow Cytometry Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.

References

1. Phillips, D.R. *et al.* (1991) GPIIb-IIIa: the responsive integrin. [Cell. 65 \(3\): 359-62.](#)
2. Michelson, A. D. *et al.* (1995) A panel of platelet mAb for the study of haemostasis and thrombosis in baboons. Leucocyte Typing V. Oxford University Press p 1230-1.
3. Comazzi, S. *et al.* (2010) Acute megakaryoblastic leukemia in dogs: a report of three cases and review of the literature. [J Am Anim Hosp Assoc. 46 \(5\): 327-35.](#)
4. Tehrani, S. *et al.* (2010) Atorvastatin has antithrombotic effects in patients with type 1 diabetes and dyslipidemia. [Thromb Res. 126 \(3\): e225-31.](#)
5. Dircks, B.H. *et al.* (2012) Platelet-neutrophil aggregate formation in blood samples from dogs with systemic inflammatory disorders. [Am J Vet Res. 73 \(7\): 939-45.](#)
6. Kamat, V. *et al.* (2015) Microfluidic assessment of functional culture-derived platelets in human thrombi under flow. [Exp Hematol. 43 \(10\): 891-900.e4.](#)
7. Ogino, T. *et al.* (2014) Erythroid and megakaryocytic differentiation of K562 erythroleukemic cells by monochloramine. [Free Radic Res. 48 \(3\): 292-302.](#)
8. Jackson, J.W. *et al.* (2016) Novel Antiplatelet Activity of Minocycline Involves Inhibition of MLK3-p38 Mitogen Activated Protein Kinase Axis. [PLoS One. 11 \(6\): e0157115.](#)
9. Chen, C.P. *et al.* (2018) Oxidative stress reduces trophoblast FOXO1 and integrin $\beta 3$

expression that inhibits cell motility. [Free Radic Biol Med. 124: 189-198.](#)

Further Reading	1. Hynes, R.O. (1992) Integrins: versatility, modulation, and signaling in cell adhesion. Cell. 69 (1): 11-25.
------------------------	--

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
----------------	--

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2588GA 10040
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Goat Anti Mouse IgG (STAR70...)	FITC

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M383948:210513'

Printed on 18 Jan 2024