

Datasheet: MCA2607 BATCH NUMBER 162223

| Description: | MOUSE ANTI HUMAN iC3b (NEOANTIGEN) |
|---------------|-------------------------------------|
| Specificity: | iC3b (NEOANTIGEN) |
| Other names: | INACTIVATED COMPLEMENT COMPONENT 3b |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | 013III-1.16 |
| Isotype: | lgG2b |
| Quantity: | 0.1 mg |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . | | | | | |
|-----------------|--|------------|-----------|--------------------|--------------------|--|
| | | Yes | No | Not Determined | Suggested Dilution | |
| | Flow Cytometry | • | | | | |
| | Immunohistology - Frozen | | | | | |
| | Immunohistology - Paraffin | • | | | 10 ug/ml | |
| | ELISA | • | | | | |
| | Immunoprecipitation | | | | | |
| | Western Blotting | • | | | | |
| | Functional Assays | | | | | |
| | Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. | | | | | |
| Target Species | Human | | | | | |
| Product Form | Purified IgG - liquid | | | | | |
| Preparation | Purified IgG prepared by | affinity c | hromatogi | raphy on Protein A | | |
| Buffer Solution | Borate buffered saline | | | | | |

| Preservative | <0.1% Sodium Azide (NaN ₃) |
|--------------|--|
| Stabilisers | |

| Approx. Protein Concentrations | Current, batch-specific concentration 1.045 mg/ml |
|--------------------------------------|---|
| Immunogen | Purified human iC3b |
| External Database Links | UniProt: P01024 Related reagents Entrez Gene: 718 C3 Related reagents |
| Synonyms | CPAMD1 |
| RRID | AB_877464 |
| Specificity | Mouse anti Human iC3b (Neoantigen) antibody, clone 013III-1.16 recognizes the ~42 kDa human inactive complement component 3b (iC3b) neoantigen present in blood serum. Cleavage of C3b into iC3b by the protease factor I, in the presence of cofactors, changes the structure and resulting binding properties, preventing binding by complement factor B and properdin. This prevents further progression of the complement pathway, providing another mechanism by which the complement pathway may be regulated. iC3b is thought to play a role in inducing tolerance (Sohn <i>et al.</i> 2003), binding complement receptor type 3 on antigen presenting cells and stimulating them to produce transforming growth factor beta2 and interleukin-10. It is thought to stimulate B-cells via the CD21 receptor providing a link between the innate and adaptive immune responses. Additionally iC3b acts as an opsonin which is recognised by complement receptor 3 expressed on leukocytes or complement receptor of the tagged cells. iC3b levels are often elevated in diseases such as Systemic Lupus Erythematosis and Rheumatoid arthritis. |
| Histology Positive Control Tissue | Kidney from post streptoccal glomerulonephritis patients |
| Further Reading | Janssena, B. <i>et al.</i> (2007) Structural insights into the central complement component C3. <u>Molecular Immunology. 44: 3-10.</u> Bergmann-Leitner, E.S. <i>et al.</i> (2006) Complement 3d: from molecular adjuvant to target of immune escape mechanisms. <u>Clin Immunol. 121 (2): 177-85.</u> Sohn, J.H. <i>et al.</i> (2003) Tolerance is dependent on complement C3 fragment iC3b binding to antigen-presenting cells. <u>Nat Med. 9 (2): 206-12.</u> |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. |
| | Avoid repeated freezing and thawing as this may denature the antibody. Storage in |

frost-free freezers is not recommended.

| Guarantee | Guaranteed until date of expiry. Please see product label. |
|----------------------------------|---|
| Health And Safety Information | Material Safety Datasheet documentation #10077 available at: https://www.bio-rad-antibodies.com/SDS/MCA2607 10077 |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

| Rabbit Anti Mouse IgG (STAR12) RPE | | | | | |
|--|-----|--|--|--|--|
| Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u> | | | | | |
| Goat Anti Mouse IgG (STAR76) RPE | | | | | |
| Rabbit Anti Mouse IgG (STAR13) <u>HRP</u> | | | | | |
| Goat Anti Mouse IgG (STAR70) <u>FITC</u> | | | | | |
| Goat Anti Mouse IgG (H/L) (STAR117) <u>Alk. Phos.</u> , <u>DyLight®488, DyLight®550</u> , | | | | | |
| DyLight®650, DyLight®680, DyLight®800, | | | | | |
| <u>FITC</u> , <u>HRP</u> | | | | | |
| Rabbit Anti Mouse IgG (STAR9) <u>FITC</u> | | | | | |
| | | | | | |
| Goat Anti Mouse IgG (STAR77) <u>HRP</u> | | | | | |
| Goat Anti Mouse IgG (Fc) (STAR120) <u>FITC</u> , <u>HRP</u> | | | | | |
| Recommended Negative Controls | | | | | |
| MOUSE IgG2b NEGATIVE CONTROL (MCA691) | | | | | |
| MOUSE IGOZD NEGATIVE CONTROL (MCAUST) | | | | | |
| North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21 | | | | | |
| America Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 | | | | | |
| Email: antibody_sales_us@bio-rad.com Email: antibody_sales_uk@bio-rad.com Email: antibody_sales_de@bio-rad.com | om | | | | |
| To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets | | | | | |
| 'M403825:220812' | 013 | | | | |

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint