

Datasheet: MCA2608

Description:	MOUSE ANTI HUMAN C4c
Specificity:	C4c
Other names:	COMPLEMENT COMPONENT 4c
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	033D-69.4.1 (10-12)
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
ELISA (1)	▪			
Western Blotting	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Clone 10-12 binds both C4c and C4 and so is not suitable for all ELISA applications.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from ascites
Buffer Solution	Borate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein	Current, batch-specific concentration 1.15 mg/ml

Concentrations

Immunogen Purified human C4c.

External Database Links

UniProt:

[P0C0L5](#) [Related reagents](#)

Entrez Gene:

[721](#) C4B [Related reagents](#)

Synonyms CO4, CPAMD3

RRID AB_844529

Specificity

Mouse anti Human C4c antibody, clone 033D-69.4.1 (10-12) recognizes complement component 4c (C4c). The complement 1 complex cleaves complement 4 (C4) to form C4b and C4a. C4b levels are strictly regulated. Single site cleavage of the C4b's alpha chain by Factor I forms iC4b and blocks C3 convertase, inhibiting opsonisation and activation of the classical pathway. iC4b is further degraded into the inactive fragments C4d and C4c. C4c consists of two a-chain fragments, alpha 4 (13 kDa) and alpha 3 (25 kDa) linked to a beta chain through disulphide bridges and a gamma chain. It is expressed in the blood plasma.

Histology Positive Control Tissue

Kidney sections from post-streptococcal glomerulonephritis patients.

Further Reading

1. Press, E.M. & Gagnon, J. (1981) Human complement component C4. Structural studies on the fragments derived from C4b by cleavage with C3b inactivator. [Biochem J. 199 \(2\): 351-7.](#)
 2. Leung, E. *et al.* (2006) The complement regulator C4b-binding protein (C4BP) interacts with both the C4c and C4dg subfragments of the parent C4b ligand: evidence for synergy in C4BP subsite binding. [Biochemistry. 45 \(27\): 8378-92.](#)
-

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

Guaranteed until date of expiry. Please see product label.

Health And Safety Information

Material Safety Datasheet documentation #10077 available at: 10077: <https://www.bio-rad-antibodies.com/uploads/MSDS/10077.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M405863:220916'

Printed on 16 Sep 2022