

## Datasheet: MCA2612

<b>Description:</b>	MOUSE ANTI HUMAN CLUSTERIN
<b>Specificity:</b>	CLUSTERIN
<b>Other names:</b>	APOLIPOPROTEIN J
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	3R3/2
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting	▪			
Functional Assays			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A
<b>Buffer Solution</b>	Borate buffered saline
<b>Preservative Stabilisers</b>	0.1% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Glomerular basement membrane from a post-mortem kidney.
<b>External Database</b>	<b>UniProt:</b>

Links [P10909](#) [Related reagents](#)

**Entrez Gene:**

[1191](#) CLU [Related reagents](#)

---

**Synonyms** APOJ, CLI, KUB1

---

**RRID** AB\_2260877

---

**Specificity** **Mouse anti Human clusterin antibody, clone 3R3/2** recognizes clusterin, an ubiquitously expressed protein of two 40 kDa chains, alpha and beta, covalently joined by disulfide bonds. It exists in two forms; a pro-apoptotic form localised to the nucleus and a pro-survival secreted form. Both are thought to be involved in DNA repair and cell cycle regulation. The secreted form is thought to inhibit apoptotic signalling by BAX, Clusterin suppression can sensitise cells to chemotherapy. Clusterin is overexpressed in human prostate cancer, breast cancers and squamous cell carcinomas.

Part of the SC5b-9 complex, clusterin prevents the binding of a C5b-C7 complex to the membrane of the target cell inhibiting the complement cascade. Clusterin acts as an extracellular chaperone inhibiting the formation of human lysozyme amyloid associated with Alzheimer's disease. Clusterin is also important in spermatogenesis and is secreted by Sertoli cells.

---

**Histology Positive Control Tissue** Kidney from post streptococcal glomerulonephritis patients.

---

**References**

1. Kumita, J.R. *et al.* (2007) The extracellular chaperone clusterin potently inhibits human lysozyme amyloid formation by interacting with prefibrillar species. [J Mol Biol. 369 \(1\): 157-67.](#)
2. Shannan, B. *et al.* (2006) Clusterin and DNA repair: a new function in cancer for a key player in apoptosis and cell cycle control. [J Mol Histol. 37 \(5-7\): 183-8.](#)

---

**Storage** Store at +4°C or at -20°C if preferred.  
Storage in frost-free freezers is not recommended.  
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** Guaranteed until date of expiry. Please see product label.

---

**Health And Safety Information** Material Safety Datasheet documentation #10334 available at:  
10334: <https://www.bio-rad-antibodies.com/uploads/MSDS/10334.pdf>

---

**Regulatory** For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)  
Goat Anti Mouse IgG (STAR77...) [HRP](#)  
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)  
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@680](#),  
[DyLight@800](#), [FITC](#), [HRP](#)

## Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M373084:200819'

Printed on 19 Aug 2020

---

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)