

Datasheet: MCA2639

BATCH NUMBER 157268

Description:	MOUSE ANTI HUMAN CYSTATIN C		
Specificity:	CYSTATIN C		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	Cyst24		
Isotype:	lgG1		
Quantity:	0.2 mg		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA	-			
Immunoprecipitation				
Western Blotting				
Immunoassay	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	A from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)	

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: P01034 Related reagents
	Entrez Gene: 1471 CST3 Related reagents
RRID	AB_2088051
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the Sp2/0 myeloma cell line.
Specificity	Mouse anti human Cystatin C antibody, clone Cyst24 detects human Cystatin C, a member of the type 2 cystatin superfamily. Cystatin C is a potent inhibitor of cysteine proteinases, both endogenous (lysosomal cathepsins), and pathogenic. Cystatin C is a ~13 kDa secreted protein which is found in high concentrations in biological fluids and is thought to be expressed by all human nucleated cells.
	Changes in cystatin C have been documented in a number of cardiovascular diseases, including atherosclerosis, amyloid angiopathy and myocardial infarction. Its low molecular weight also means that cystatin C can be used as a marker of renal failure.
References	1. Mussap, M. & Plebani, M. (2004) Biochemistry and clinical role of human cystatin C. Crit Rev Clin Lab Sci. 41 (5-6): 467-550.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2639 10040

Related Products

Regulatory

Recommended Secondary Antibodies

For research purposes only

Rabbit Anti Mouse IgG (STAR12...)

RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)

RPE
Goat Anti Mouse IgG (STAR76...)

RPE

Rabbit Anti Mouse IgG (STAR13...) HRP
Goat Anti Mouse IgG (STAR70...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M378861:210309'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint