

# Datasheet: MCA2639 BATCH NUMBER 161666

Description:	MOUSE ANTI HUMAN CYSTATIN C		
Specificity:	CYSTATIN C		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	Cyst24		
lsotype:	lgG1		
Quantity:	0.2 mg		

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further						
	information. For general p	information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry			•			
	Immunohistology - Frozen						
	Immunohistology - Paraffin						
	ELISA	•					
	Immunoprecipitation						
	Western Blotting						
	Immunoassay						
	Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as						
	a guide only. It is recommended that the user titrates the product for use in their own						
	system using appropriate negative/positive controls.						
Target Species	Human						
Product Form	Purified IgG - liquid						
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant						
Buffer Solution	Phosphate buffered salin	e					
Preservative Stabilisers	<0.1% Sodium Azide (Na	N <sub>3</sub> )					

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt: <u>P01034</u> <u>Related reagents</u> Entrez Gene: <u>1471</u> CST3 <u>Related reagents</u>
RRID	AB_2088051
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the Sp2/0 myeloma cell line.
Specificity	Mouse anti human Cystatin C antibody, clone Cyst24 detects human Cystatin C, a member of the type 2 cystatin superfamily. Cystatin C is a potent inhibitor of cysteine proteinases, both endogenous (lysosomal cathepsins), and pathogenic. Cystatin C is a ~13 kDa secreted protein which is found in high concentrations in biological fluids and is thought to be expressed by all human nucleated cells. Changes in cystatin C have been documented in a number of cardiovascular diseases, including atherosclerosis, amyloid angiopathy and myocardial infarction. Its low molecular weight also means that cystatin C can be used as a marker of renal failure.
References	1. Mussap, M. & Plebani, M. (2004) Biochemistry and clinical role of human cystatin C. <u>Crit Rev Clin Lab Sci. 41 (5-6): 467-550.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA2639 10040
Regulatory	For research purposes only

## **Related Products**

### **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (STAR77)	<u>HRP</u>
Rabbit Anti Mouse IgG (STAR12)	<u>RPE</u>

Rabbit Anti Mouse IgG (STAR8)	DyLight®800
Goat Anti Mouse IgG (STAR76)	RPE
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP
Goat Anti Mouse IgG IgA IgM (STAR87	) <u>Alk. Phos.</u> , <u>HRP</u>
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,
	DyLight®650, DyLight®680, DyLight®800,
	<u>FITC, HRP</u>
<b>Decommonded Negative Controls</b>	

#### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M389447:210806'

#### Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint