Datasheet: MCA2648 BATCH NUMBER 167522

| Description: | MOUSE ANTI HUMAN C3d |
|---------------|-------------------------|
| Specificity: | C3d |
| Other names: | COMPLEMENT COMPONENT 3d |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | 053A-514.3.1.4 |
| Isotype: | lgG1 |
| Quantity: | 0.1 mg |
| | |

Product Details

 Applications
 This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.biorad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--|-------------------------|----------------------|--|-------------------------|
| Flow Cytometry | | | | |
| Immunohistology - Frozen | • | | | |
| Immunohistology - Paraffin | | | • | |
| ELISA | | | | |
| Immunoprecipitation | | | • | |
| Western Blotting | | | • | |
| Functional Assays | | | • | |
| Where this product has n necessarily exclude its us a guide only. It is recomm system using appropriate | se in such nended th | procedu at the us | ures. Suggested workin ser titrates the product f | g dilutions are given a |

| Target Species | Human | |
|-----------------------------|---|--------------|
| Product Form | Purified IgG - liquid | |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A | from ascites |
| Buffer Solution | Borate buffered saline | |
| Preservative Stabilisers | 0.1% Sodium Azide (NaN ₃) | |

| Approx. Protein Concentrations | Current, batch-specific concentration 1.125 mg/ml |
|--------------------------------------|---|
| Immunogen | Purified human C3d (neoantigen). |
| External Database Links | UniProt: P01024 Related reagents Entrez Gene: 718 C3 Related reagents |
| Synonyms | CPAMD1 |
| RRID | AB_2259274 |
| Specificity | Mouse anti Human C3d antibody, clone 053A-514.3.1.4 recognizes human complement component 3d (C3d) neoantigen, a ~33 kDa polypeptide fragment generated over the course of complement activation in which where complement 3 (C3) convertases cleave C3 to C3b, which is further degraded into iC3b and C3dg/C3d. C3d is involved in the regulation of many aspects of the immune response, including antigen processing and presentation. The C3d/antigen complex mediates B cell activation by simultaneously binding antigen-specific surface bound immunoglobulin and CD21, lowering the complement activation threshold. C3 is crucial in the induction of tolerance generated when an antigen is introduced into immunoprivileged sites and this is exploited by pathogens and cancer cells to evade the immune system by inhibiting complement activation. |
| Histology Positive Control Tissue | Kidney from post streptococcal glomerulonephritis patients |
| References | 1. Verghese, P.S. <i>et al.</i> (2018) The clinical implications of the unique glomerular complement deposition pattern in transplant glomerulopathy. <u>J Nephrol. 31 (1): 157-64.</u> |
| Further Reading | Bergmann-Leitner, E.S. <i>et al.</i> (2006) Complement 3d: from molecular adjuvant to target of immune escape mechanisms. <u>Clin Immunol. 121 (2): 177-85.</u> Barilla-LaBarca, M.L. <i>et al.</i> (2002) Role of membrane cofactor protein (CD46) in regulation of C4b and C3b deposited on cells. <u>J Immunol. 168 (12): 6298-304.</u> |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. |
| | Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. |
| Guarantee | Guaranteed until date of expiry. Please see product label. |

| Health And Safety | Material Safety Datasheet documentation #10077 available at: | | | |
|-------------------|--|--|--|--|
| Information | https://www.bio-rad-antibodies.com/SDS/MCA2648 | | | |
| | 10077 | | | |
| | | | | |

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

| Rabbit Anti Mouse IgG (STAR12) | RPE | | |
|---|---|--|--|
| Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u> | | | |
| Goat Anti Mouse IgG (STAR76) | RPE | | |
| Goat Anti Mouse IgG (STAR70) | <u>FITC</u> | | |
| Goat Anti Mouse IgG (H/L) (STAR117) | <u>Alk. Phos.</u> , <u>DyLight®488</u> , <u>DyLight®550</u> , | | |
| | DyLight®650, DyLight®680, DyLight®800, | | |
| | <u>FITC</u> , <u>HRP</u> | | |
| Rabbit Anti Mouse IgG (STAR9) | <u>FITC</u> | | |
| Goat Anti Mouse IgG (STAR77) | HRP | | |
| Goat Anti Mouse IgG (Fc) (STAR120) | FITC, HRP | | |
| Rabbit Anti Mouse IgG (STAR13) | HRP | | |
| Recommended Negative Controls | | | |
| | | | |

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 |
|---------------|---------------------------------|-----------|-------------------------------|---------|--------------------------------------|
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 |
| | Email: antibody_sales_us@bio-ra | id.com | Email: antibody_sales_uk@bio- | rad.com | Email: antibody_sales_de@bio-rad.com |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M426641:240212'

Printed on 12 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint