

## Datasheet: MCA2659

<b>Description:</b>	MOUSE ANTI HUMAN CD55
<b>Specificity:</b>	CD55
<b>Other names:</b>	DAF
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	Bu14
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

**RRID** AB\_2075955

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Western Blotting			▪	
Functional Assays			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

**Target Species** Human

**Product Form** Purified IgG - liquid

**Preparation** Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

**Buffer Solution** Phosphate buffered saline

**Preservative Stabilisers** 0.09% Sodium Azide (NaN<sub>3</sub>)

**Carrier Free** Yes

**Approx. Protein Concentrations** IgG concentration 1.0mg/ml

**External Database  
Links**

**UniProt:**

[P08174](#)   [Related reagents](#)

**Entrez Gene:**

[1604](#)   CD55   [Related reagents](#)

---

**Synonyms**

CR, DAF

---

**Specificity**

**Mouse anti Human CD55 antibody, clone Bu14** recognizes human CD55, also known as Decay Accelerating Factor (DAF). CD55 is a glycosylphosphatidylinositol (GPI) anchored type I surface protein which prevents complement mediated lysis of autologous cells. CD55 functions to limit the formation and accelerate the decay of C3/C5-convertases from the classic and alternative pathway.

The molecular weight of CD55 varies between ~50 and 100 kDa dependent on cell type. Alternative transcripts can lead to soluble DAF, which is detectable in plasma, tears, saliva, urine and cerebrospinal fluid ([Medof et al. 1987](#)).

CD55 acts as a ligand for CD97 ([Hamaan et al. 1998](#)) and is an entry receptor for Echovirus ([Bergelson et al. 1994](#)) and Coxsackie B virus ([Hafenstein et al. 2007](#)). CD55 has been linked to many diseases including multiple sclerosis, rheumatoid arthritis and cancer development ([Visser et al. 2002](#)).

---

**Flow Cytometry**

Use 10ul of the suggested working dilution to label 1x10<sup>6</sup> cells in 100ul.

---

**References**

1. Loveland, B.E. and Szokolai, K. (1998) CD55 Workshop Report NL11 In: Leukocyte Typing VI, Eds Kishimoto, T. et al. Garland publishing . 519-520.

---

**Further Reading**

1. Ruiz-Argüelles, A. & Llorente, L. (2007) The role of complement regulatory proteins (CD55 and CD59) in the pathogenesis of autoimmune hemocytopenias. [Autoimmun Rev. 6 \(3\): 155-61.](#)  
2. Mikesch, J.H. et al. (2006) Decay-accelerating factor (CD55): a versatile acting molecule in human malignancies. [Biochim Biophys Acta. 1766 \(1\): 42-52.](#)

---

**Storage**

Store at +4°C or at -20°C if preferred.  
Storage in frost-free freezers is not recommended.  
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

18 months from date of despatch.

---

**Health And Safety  
Information**

Material Safety Datasheet documentation #10040 available at:  
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)

Goat Anti Mouse IgG (STAR77...) [HRP](#)

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)

Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®680](#),  
[DyLight®800](#), [FITC](#), [HRP](#)

## Recommended Negative Controls

### [MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M343471:190111'

Printed on 20 May 2019

---

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)