

Datasheet: MCA2659F

Description:	MOUSE ANTI HUMAN CD55:FITC
Specificity:	CD55
Other names:	DAF
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	Bu14
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

RRID AB_2275855

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/20

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Human

Product Form Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative 0.09% Sodium Azide (NaN₃)
Stabilisers 1% Bovine Serum Albumin

Approx. Protein Concentrations IgG concentration 0.1 mg/ml

External Database Links
UniProt:
[P08174](#) [Related reagents](#)

Entrez Gene:

[1604](#) CD55 [Related reagents](#)

Synonyms	CR, DAF
-----------------	---------

Specificity	<p>Mouse anti Human CD55 antibody, clone Bu14 recognizes human CD55, also known as Decay Accelerating Factor (DAF). CD55 is a glycosylphosphatidylinositol (GPI) anchored type I surface protein which prevents complement mediated lysis of autologous cells. CD55 functions to limit the formation and accelerate the decay of C3/C5-convertases from the classic and alternative pathway.</p> <p>The molecular weight of CD55 varies between ~50 and 100 kDa dependent on cell type. Alternative transcripts can lead to soluble DAF, which is detectable in plasma, tears, saliva, urine and cerebrospinal fluid (Medof et al. 1987).</p> <p>CD55 acts as a ligand for CD97 (Hamaan et al. 1998) and is an entry receptor for Echovirus (Bergelson et al. 1994) and Coxsackie B virus (Hafenstein et al. 2007). CD55 has been linked to many diseases including multiple sclerosis, rheumatoid arthritis and cancer development (Visser et al. 2002).</p>
--------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
-----------------------	-------------------------------------------------------------------------------------

References	1. Loveland, B.E. and Szokolai, K. (1998) CD55 Workshop Report NL11 In: Leukocyte Typing VI, Eds Kishimoto, T. et al. Garland publishing . 519-520.
-------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

Further Reading	1. Ruiz-Argüelles, A. & Llorente, L. (2007) The role of complement regulatory proteins (CD55 and CD59) in the pathogenesis of autoimmune hemocytopenias. Autoimmun Rev. 6 (3): 155-61. 2. Mikesch, J.H. et al. (2006) Decay-accelerating factor (CD55): a versatile acting molecule in human malignancies. Biochim Biophys Acta. 1766 (1): 42-52.
------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
----------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Guarantee	18 months from date of despatch.
------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
--------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com
'M348259:190219'

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)