Datasheet: MCA2674PE BATCH NUMBER 153612

Description:	HAMSTER ANTI HUMAN EMR1:RPE
Specificity:	EMR1
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	A10
Isotype:	lgG
Quantity:	100 TESTS

Product Details

Applications	This product has been r derived from testing with communications from th information. For general rad-antibodies.com/prote	nin our labo e originato protocol re	oratories, rs. Pleas	peer-reviewed publicate e refer to references in	tions or personal dicated for further	
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry				Neat	
	Where this product has not been tested for use in a particular technique this does non- necessarily exclude its use in such procedures. Suggested working dilutions are giv a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.					
Target Species	Human					
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized					
Reconstitution	Reconstitute with 1.0ml distilled water					
Max Ex/Em	Fluorophore	Excitation N	lax (nm)	Emission Max (nm)		
	RPE 488nm laser	496		578		
Preparation	Purified IgG prepared by affinity chromatography on Protein G					
Buffer Solution	Phosphate buffered saline					
Preservative	0.09% Sodium Azide					
Stabilisers	1% Bovine Serum Albumin					
	5% Sucrose					

External Database Links UniProt: Q14246 Related reagents Entrez Gene: 2015 EMR1 Related reagents Synonyms TM7LN3 RRID AB_1085232 Fusion Partners Spleen cells from immunised Armenian hamster were fused with cells of the mouse SP2/0 myeloma cell line. Specificity Hamster anti Human EMR1 antibody, clone A10 recognizes human EMR1 (EGF-like module containing, mucin-like, hormone receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/80 but unlike F4/80, the expression of EMR1 in humans is restricted to eosinophils and is not expressed on macrophage populations. The detailed structure and function of EMR1 is still unknown. Flow Cytometry Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. References 1. van Eijk, M. <i>et al.</i> (2010) Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. Immunol Lett. 129 (2): 64-71. 2. Pothoven KL <i>et al.</i> (2011) Associations Armong Adipose Tissue Immunology, Inflammation, Exosomes and Insulin Sensitivity in People With Obesity and Nonalcoholic Fatty Liver Disease. Gastroenterology. 161 (3): 968-981.e12. Storege Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. <td< th=""><th></th><th></th></td<>		
Q14246 Related reagents Entrez Gene: 2015 2015 EMR1 RRID AB_10852232 Fusion Partners Spleen cells from immunised Armenian hamster were fused with cells of the mouse SP2/0 myeloma cell line. Specificity Hamster anti Human EMR1 antibody, clone A10 recognizes human EMR1 (EGF-like module containing, mucin-like, hormone receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/40, the expression of EMR1 in humans is restricted to eosinophils and is not expressed on macrophage populations. The detailed structure and function of EMR1 is still unknown. Flow Cytometry Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. References 1. van Eijk, M. <i>et al.</i> (2010) Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atheroscierosis, multiple sclerosis and Gaucher disease. Immunol Lett. 129 (2): 64-71. .2. Pothoven KL <i>et al.</i> (2010) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J. Allergy Clin Immunol. pii: S0091-6749(16)31462-2. Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Materia		UniProt:
2015 EMR1 Related reagents Synonyms TM7LN3 RRID AB_10852232 Fusion Partners Spleen cells from immunised Armenian hamster were fused with cells of the mouse SP2/0 myeloma cell line. Specificity Hamster anti Human EMR1 antibody, clone A10 recognizes human EMR1 (EGF-like module containing, mucin-like, hormone receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/80 but unlike F4/80, the expression of EMR1 in humans is restricted to eosinophils and is not expression of EMR1 in humans is restricted to eosinophils and is not expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. Immunol. Ett. 129 (2): 64-71. References 1. van Eijk, M. <i>et al.</i> (2010) Nettrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J. Altergy Clin Immunol. pii: S0091-6749(16)31462-2. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDSMCA2674PE	LIIKS	
Synonyms TM7LN3 RRID AB_10852232 Fusion Partners Spleen cells from immunised Armenian hamster were fused with cells of the mouse SP2/0 myeloma cell line. Specificity Hamster anti Human EMR1 antibody, clone A10 recognizes human EMR1 (EGF-like module containing, mucin-like, hormore receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/80 but unlike F4/80, the expression of EMR1 in humans is restricted to eosinophils and is not expressed on macrophage populations. The detailed structure and function of EMR1 is still unknown. Flow Cytometry Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. References 1. van Eijk, M. <i>et al.</i> (2010) Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher discase. Immunol Lett. 129 (2): 64-71. 2. Pothoven KL <i>et al.</i> (2016) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J Allergy Clin Immunol. pit: S0091-6749(16):1462-22. 3. Fuchs, A. <i>et al.</i> (2021) Associations Among Adipose Tissue Immunology, Inflammation, Exosomes and Insulin Sensitivity in People With Obesity and Nonalcoholic Fatty Liver Disease. Gastroenterology. 161 (3): 968-981.e12. Storege Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.		Entrez Gene:
RRID AB_10852232 Fusion Partners Spleen cells from immunised Armenian hamster were fused with cells of the mouse SP2/0 myeloma cell line. Specificity Hamster anti Human EMR1 antibody, clone A10 recognizes human EMR1 (EGF-like module containing, mucin-like, hormone receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/80 but unlike F4/80, the expression of EMR1 in humans is restricted to eosinophils and is not expressed on macrophage populations. The detailed structure and function of EMR1 is still unknown. Flow Cytometry Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. References 1. van Eijk, M. <i>et al.</i> (2010) Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. Immunol Lett. 129 (2): 64-71. 2. Pothoven KL <i>et al.</i> (2016) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J Allergy Clin Immunol. pii: S0091-6749(16)31462-2. 3. Fuchs, A. <i>et al.</i> (2021) Associations Among Adipose Tissue Immunology, Inflammation, Exosomes and Insulin Sensitivity in People With Obesity and Nonalcoholic Fatty Liver Disease. Gastroenterology, 161 (3): 968-981.e12. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undituded. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee		2015 EMR1 Related reagents
Fusion Partners Spieen cells from immunised Armenian hamster were fused with cells of the mouse SP2/0 myeloma cell line. Specificity Hamster anti Human EMR1 antibody, clone A10 recognizes human EMR1 (EGF-like module containing, mucin-like, hormone receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/80 but unlike F4/80, the expression of EMR1 in humans is restricted to eosinophilis and is not expressed on macrophage populations. The detailed structure and function of EMR1 is still unknown. Flow Cytometry Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. References 1. van Eijk, M. <i>et al.</i> (2010) Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. Immunol Lett. 129 (2): 64-71. 2. Pothoven KL <i>et al.</i> (2010) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J Allergy Clin Immunol. pli: S0091-6749(16)31462-2. 3. Fuchs, A. <i>et al.</i> (2021) Associations Among Adipose Tissue Immunology. Inflammation, Exosomes and Insultin Sensitivity in People With Obesity and Nonalcoholic Fatty Liver Disease. Gastroenterology, 161 (3): 968-981.e12. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch	Synonyms	TM7LN3
Fusion Partners Spleen cells from immunised Armenian hamster were fused with cells of the mouse SP2/0 myeloma cell line. Specificity Hamster anti Human EMR1 antibody, clone A10 recognizes human EMR1 (EGF-like module containing, mucin-like, hormone receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/80 but unlike F4/80, the expression of EMR1 in humans is restricted to eosinophils and is not expressed on macrophage populations. The detailed structure and function of EMR1 is still unknown. Flow Cytometry Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. References 1. van Eijk, M. <i>et al.</i> (2010) Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosolerosis, multiple sclerosis and Gaucher disease. Immunol Lett. 129 (2): 64-71. 2. Pothoven KL <i>et al.</i> (2010) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J Allergy Clin Immunol. <i>pli:</i> S0091-6749(16)31462-2. 3. Fuchs, A. <i>et al.</i> (2021) Associations Among Adipose Tissue Immunology. Inflammation, Exosomes and Insulin Sensitivity in People With Obesity and Nonalcoholic Fatty Liver Disease. Gastroenterology. 161 (3): 968-981.e12. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch	RRID	AB 10852232
Specificity Hamster anti Human EMR1 antibody, clone A10 recognizes human EMR1 (EGF-like module containing, mucin-like, hormone receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/80 but unlike F4/80, the expression of EMR1 in humans is restricted to eosinophils and is not expressed on macrophage populations. The detailed structure and function of EMR1 is still unknown. Flow Cytometry Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. References 1. van Eijk, M. <i>et al.</i> (2010) Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. Immunol Lett. 129 (2): 64-71. 2. Pothoven KL <i>et al.</i> (2010) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J Allergy Clin Immunol. pii: S0091-6749(16)31462-2. 3. Fuchs, A. <i>et al.</i> (2021) Associations Among Adipose Tissue Immunology, Inflammation, Exosomes and Insulin Sensitivity in People With Obesity and Nonalcoholic Fatty Liver Disease. Gastroenterology. 161 (3): 968-981.e12. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2674PE 20487		_
module containing, mucin-like, hormone receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/80 but unlike F4/80, the expression of EMR1 in humans is restricted to eosinophils and is not expressed on macrophage populations. The detailed structure and function of EMR1 is still unknown. Flow Cytometry Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul. References 1. van Eijk, M. <i>et al.</i> (2010) Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. Immunol Lett. 129 (2): 64-71. 2. Pothoven KL <i>et al.</i> (2016) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J Allergy Clin Immunol. pii: S0091-6749(16)31462-2. 3. Fuchs, A. <i>et al.</i> (2021) Associations Among Adipose Tissue Immunology, Inflammation, Exosomes and Insulin Sensitivity in People With Obesity and Nonalcoholic Fatty Liver Disease. Gastroenterology, 161 (3): 968-981.e12. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2674PE 20487	Fusion Partners	
References 1. van Eijk, M. et al. (2010) Differential expression of the EGF-TM7 family members CD97 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. Immunol Lett. 129 (2): 64-71. 2. Pothoven KL et al. (2016) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J Allergy Clin Immunol. pii: S0091-6749(16)31462-2. 3. Fuchs, A. et al. (2021) Associations Among Adipose Tissue Immunology, Inflammation, Exosomes and Insulin Sensitivity in People With Obesity and Nonalcoholic Fatty Liver Disease. Gastroenterology. 161 (3): 968-981.e12. Storage Store at +4°C. DO NOT FREEZE. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2674PE 20487	Specificity	module containing, mucin-like, hormone receptor-like 1). EMR1 is a seven-span transmembrane protein which is a member of the EGF-TM7 family of receptors. EMR1 is the human ortholog of murine F4/80 but unlike F4/80, the expression of EMR1 in humans is restricted to eosinophils and is not expressed on macrophage populations. The detailed
and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. Immunol Lett. 129 (2): 64-71. 2. Pothoven KL <i>et al.</i> (2016) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. J Allergy Clin Immunol. pii: S0091-6749(16)31462-2. 3. Fuchs, A. <i>et al.</i> (2021) Associations Among Adipose Tissue Immunology, Inflammation, Exosomes and Insulin Sensitivity in People With Obesity and Nonalcoholic Fatty Liver Disease. Gastroenterology. 161 (3): 968-981.e12.StorageStore at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.Guarantee12 months from date of despatchHealth And Safety InformationMaterial Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2674PE 20487	Flow Cytometry	Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.
StorageStore at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.Guarantee12 months from date of despatchHealth And Safety InformationMaterial Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2674PE 20487	References	 and EMR2 in lipid-laden macrophages in atherosclerosis, multiple sclerosis and Gaucher disease. <u>Immunol Lett. 129 (2): 64-71.</u> 2. Pothoven KL <i>et al.</i> (2016) Neutrophils are a major source of the epithelial barrier disrupting cytokine oncostatin M in patients with mucosal airways disease. <u>J Allergy Clin Immunol. pii: S0091-6749(16)31462-2.</u> 3. Fuchs, A. <i>et al.</i> (2021) Associations Among Adipose Tissue Immunology, Inflammation,
This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #20487 available at: https://www.bio-rad-antibodies.com/SDS/MCA2674PE		Disease. Gastroenterology. 161 (3): 968-981.e12.
Health And Safety Material Safety Datasheet documentation #20487 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA2674PE 20487	Storage	This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend
Information <u>https://www.bio-rad-antibodies.com/SDS/MCA2674PE</u> 20487	Guarantee	12 months from date of despatch
Regulatory For research purposes only	-	https://www.bio-rad-antibodies.com/SDS/MCA2674PE
	Regulatory	For research purposes only

Related Products

Recommended Negative Controls

HAMSTER (ARMENIAN) IgG NEGATIVE CONTROL:RPE (MCA2356PE)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Vorldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.co	om	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375522:210104'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint