

Datasheet: MCA2691SBR775 BATCH NUMBER 64589178

Description:	RAT ANTI MOUSE CD4:StarBright Red 775
Specificity:	CD4
Other names:	L3T4 ANTIGEN, LY-4
Format:	StarBright Red 775
Product Type:	Monoclonal Antibody
Clone:	RM4-5
Isotype:	lgG2a
Quantity:	100 TESTS/0.5ml

Product Details

Applications	derived from testing wi communications from t	thin our laborato he originators. P al protocol recom	in the following applicati ries, peer-reviewed public lease refer to references mendations, please visit	cations or personal indicated for further	
		Yes N	Not Determined	Suggested Dilution	
	Flow Cytometry	•		Neat	
	necessarily exclude its	use in such proo nmended that th	e user titrates the produc	ing dilutions are given as	
Target Species	Mouse				
Product Form	Purified IgG conjugated	d to StarBright R	ed 775 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm) Emission Max (nm)		
	StarBright Red 775	653	778		
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant				
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% Sodium Azide (I	NaN ₃)			
Stabilisers	1% Bovine Serum Albu	umin			
	0.1% Pluronic F68				
	0.1% PEG 3350				

	0.05% Tween 20
Immunogen	BALB/c mouse thymocytes
External Database Links	UniProt: <u>P06332</u> <u>Related reagents</u> Entrez Gene: <u>12504</u> Cd4 <u>Related reagents</u>
Specificity	Rat anti Mouse CD4 antibody, clone RM4-5 detects mouse CD4, a 55 kDa protein also known as Ly-4 and L3T4. CD4 is a single chain transmembraneous glycoprotein which belongs to the immunoglobulin superfamily, and is primarily expressed on peripheral blood monocytes and tissue macrophages. CD4 is also expressed on a subpopulation of regulatory T cells (CD4 ⁺ CD25 ⁺), which play a key role in the maintenance of self tolerance. Rat anti Mouse CD4 antibody, clone RM4-5 can be used for <i>in vitro</i> blocking of ligand binding, as well as <i>in vitro</i> CD4+ T cell depletions.
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	 von Kutzleben, S. <i>et al.</i> (2017) Depletion of CD52-positive cells inhibits the development of central nervous system autoimmune disease, but deletes an immune- tolerance promoting CD8 T-cell population. Implications for secondary autoimmunity of alemtuzumab in multiple sclerosis. Immunology. 150 (4): 444-55. Zamudio, F. <i>et al.</i> (2020) TDP-43 mediated blood-brain barrier permeability and leukocyte infiltration promote neurodegeneration in a low-grade systemic inflammation mouse model. J Neuroinflammation. 17 (1): 283. Aloui, A. <i>et al.</i> (2023) AFM(1) Exposure in Male Balb/c Mice and Intervention Strategies Against Its Immuno-physiological toxicity using Clay Mineral and Lactic Acid Bacteria Alone or in Combination. Immunopharmacol Immunotoxicol. : 1-32. Camponeschi, C. <i>et al.</i> (2021) S100B Protein as a Therapeutic Target in Multiple Sclerosis: The S100B Inhibitor Arundic Acid Protects from Chronic Experimental Autoimmune Encephalomyelitis. Int J Mol Sci. 22 (24): 13558.
Further Reading	1. Fehérvari, Z. & Sakaguchi, S. (2004) CD4+ Tregs and immune control. <u>J Clin Invest.</u> <u>114 (9): 1209-17.</u>
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts

Health A Informati		erial Safety Datas <u>s://www.bio-rad-ar</u> 71			
Regulato	ry For				
Relate	d Products				
Recomm	nended Useful	Reagents			
	EROBLOCK FcR (E EROBLOCK FcR (E				
North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
America		us@bio-rad.com	Email: antibody sales uk@bio	o-rad.com	Email: antibody sales de@bio-rad.com

Printed on 09 Apr 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint