

# Datasheet: MCA2718 BATCH NUMBER 165381

Description:	MOUSE ANTI CHLAMYDIA LPS		
Specificity:	CHLAMYDIA LPS		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	CF 6J12		
Isotype:	IgG2a		
Quantity:	1 mg		

### **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA	•			
Western Blotting	•			
Immunofluorescence	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Bacterial
Species Cross Reactivity	Reacts with: Chlamydophila sp. <b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline.
Preservative Stabilisers	<0.1% Sodium Azide (NaN <sub>3</sub> )
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Elementary bodies from <i>C. trachomatis</i> strain SAF <sub>2</sub> .
RRID	AB_915244
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NS0/U mouse myeloma cell line.
Specificity	Mouse anti Chlamydia LPS antibody, clone CF 6J12 recognizes a genus specific epitope within <i>Chlamydia</i> lipopolysaccharide (LPS).
	LPS is a common feature of the outer envelope of gram negative bacteria, which acts as a potent endotoxin, triggering an innate immune response. Whilst the LPS of <i>Chlamydia trachomatis</i> does evoke an immune response, it displays only weak endotoxic activity when compared to that of other bacteria such as <i>Salmonella minnesota</i> or <i>Neisseria gonorrhoeae</i> (Ingalls et al. 1995).
References	<ol> <li>Thornley, M.J. et al. (1985) Properties of monoclonal antibodies to the genus-specific antigen of <i>Chlamydia</i> and their use for antigen detection by reverse passive haemagglutination. J Gen Microbiol. 131 (1): 7-15.</li> <li>Xia M et al. (2013) Immunization of <i>Chlamydia pneumoniae</i> (Cpn)-infected Apob(tm2Sgy)Ldlr(tm1Her)/J mice with a combined peptide of Cpn significantly reduces atherosclerotic lesion development. PLoS One. 8 (12): e81056.</li> <li>Campbell, S. et al. (1994) Lipopolysaccharide in cells infected by <i>Chlamydia trachomatis</i>. Microbiology.140: 1995-2002.</li> <li>Xia, M. et al. (2013) Immunization of <i>Chlamydia pneumoniae</i>. (Cpn)-infected Apob(tm2Sgy)Ldlr(tm1Her)/J mice with a combined peptide of Cpn significantly reduces atherosclerotic lesion development. PLoS One. 8 (12): e81056.</li> <li>Borth, N. et al. (2011) Functional interaction between type Ill-secreted protein IncA of <i>Chlamydophila psittaci</i>. and human G3BP1. PLoS One. 6 (1): e16692.</li> <li>Dlugosz, A. et al. (2010) <i>Chlamydia trachomatis</i>. antigens in enteroendocrine cells and macrophages of the small bowel in patients with severe irritable bowel syndrome. BMC Gastroenterol. 10: 19.</li> <li>Lantos, I. et al. (2018) <i>Chlamydia pneumoniae</i> Infection Exacerbates Atherosclerosis in ApoB100only/LDLR<sup>-/-</sup> Mouse Strain. Biomed Res Int. 2018: 8325915.</li> <li>Mosolygó, T. et al. (2019) Selenocompounds as Novel Antibacterial Agents and Bacterial Efflux Pump Inhibitors. Molecules. 24(8):1487.</li> </ol>
Further Reading	1. Ingalls, R.R. <i>et al.</i> (1995) The inflammatory cytokine response to <i>Chlamydia trachomatis</i> infection is endotoxin mediated. <u>Infect Immun. 63 (8): 3125-30.</u>

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at

-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for

short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in

frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

**Health And Safety** Information

Material Safety Datasheet documentation #10040 available at:

https://www.bio-rad-antibodies.com/SDS/MCA2718

10040

Regulatory For research purposes only

#### Related Products

## **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...) **RPE** 

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...) **RPE** 

Goat Anti Mouse IgG (STAR70...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (STAR77...) **HRP** 

FITC, HRP Goat Anti Mouse IgG (Fc) (STAR120...)

Rabbit Anti Mouse IgG (STAR13...) **HRP** Rabbit Anti Mouse IgG (STAR9...) **FITC** 

North & South Tel: +1 800 265 7376

America

Worldwide Fax: +1 919 878 3751

Email: antibody\_sales\_us@bio-rad.com

Tel: +44 (0)1865 852 700

Europe Fax: +44 (0)1865 852 739

Email: antibody\_sales\_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M389469:210806'

#### Printed on 16 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint