

Datasheet: MCA2726F

BATCH NUMBER 163353

Description:	MOUSE ANTI HUMAN CD44:FITC
Specificity:	CD44
Other names:	H-CAM, PGP-1
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	156-3C11
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Baboon, African green monkey , Cat

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Stimulated human leucocytes.
External Database Links	<p>UniProt: P16070 Related reagents</p> <p>Entrez Gene: 960 CD44 Related reagents</p>
Synonyms	LHR, MDU2, MDU3, MIC4
Specificity	<p>Mouse anti Human CD44 antibody, clone 156-3C11 recognizes human Phagocytic glycoprotein 1 also known as CD44, HCAM or CD44s. CD44 is a ~90 kDa single pass type I transmembrane glycoprotein. Various isoforms of CD44 exist due to differential expression of exon products from the membrane proximal region of the extracellular domain. Mouse anti Human CD44 antibody, clone 156-3C11 recognizes the ~90 kDa standard form lacking any of the alternative spliced products, the clone is expected to recognize all isoforms of CD44. CD44 is expressed on leucocytes, erythrocytes, white matter of the brain and some epithelial cells of the breast and small intestine. Antibodies produced by clone 156-3C11 recognise epitope 3, defined as a protease resistant epitope on the CD44 molecule (CD44 and CD45R Cluster report. In Leucocyte Typing V. White cell differentiation antigens. Eds Schlossman, S.F. <i>et al</i>).</p> <p>CD44 is a receptor for hyaluronic acid (HA) and is involved in cell-cell interactions, cell adhesion and migration (Lesley <i>et al.</i> 1990). CD44 also participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing (Shimizu <i>et al.</i> 1989). CD44 expression may be up-regulated upon some carcinomas, and it has been speculated that this may be related to metastatic potential (East and Hart 1993).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul
References	<ol style="list-style-type: none"> Denning, S.M. <i>et al.</i> (1995) CD44 and CD45R Cluster report. In Leucocyte Typing V. White cell differentiation antigens. Eds Schlossman, S.F. <i>et al.</i> Oxford University Press. Volume 2,AS10:1713 – 1719 Olsson, E. <i>et al.</i> (2011) CD44 isoforms are heterogeneously expressed in breast cancer and correlate with tumor subtypes and cancer stem cell markers. BMC Cancer. 11: 418. Alves, C.S. <i>et al.</i> (2009) Biomolecular characterization of CD44-fibrin(ogen) binding: distinct molecular requirements mediate binding of standard and variant isoforms of CD44 to immobilized fibrin(ogen). J Biol Chem. 284: 1177-89. Heidemann, F. <i>et al.</i> (2014) Selectins mediate small cell lung cancer systemic

metastasis. [PLoS One. 9\(4\):e92327.](#)

5. Zhang, D. *et al.* (2016) Screening and Identification of a Phage Display Derived Peptide That Specifically Binds to the CD44 Protein Region Encoded by Variable Exons. [J Biomol Screen. 21 \(1\): 44-53.](#)

6. Zhang, P. *et al.* (2014) CD44 variant, but not standard CD44 isoforms, mediate disassembly of endothelial VE-cadherin junction on metastatic melanoma cells. [FEBS Lett. 588 \(24\): 4573-82.](#)

7. Pinto, F. *et al.* (2014) T-box transcription factor brachyury is associated with prostate cancer progression and aggressiveness. [Clin Cancer Res. 20 \(18\): 4949-61.](#)

8. Afonso, J. *et al.* (2015) CD147 and MCT1-potential partners in bladder cancer aggressiveness and cisplatin resistance. [Mol Carcinog. 54 \(11\): 1451-66.](#)

9. Lawson, J.S. *et al.* (2018) Characterisation of feline renal cortical fibroblast cultures and their transcriptional response to transforming growth factor β 1. [BMC Vet Res. 14 \(1\): 76.](#)

10. Lawson, J.S. *et al.* (2019) Characterisation of Crandell-Rees Feline Kidney (CRFK) cells as mesenchymal in phenotype. [Res Vet Sci. 127: 99-102.](#)

11. Lara, M.L. *et al.* (2023) Influence of culture conditions on the secretome of mesenchymal stem cells derived from feline adipose tissue: Proteomics approach. [Biochimie. 211: 78-86.](#)

12. Tiraihi, T. *et al.* (2023) A Sequential Culturing System for Generating Epithelial-Like Stem Cells from Human Mesenchymal Stem Cells Derived from Adipose Tissue [Cell Tiss Biol. 17 \(6\): 639-52.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA2726F10041>

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:FITC \(MCA929F\)](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M418966:230502'

