

Datasheet: MCA2733GA

Description:	MOUSE ANTI HUMAN MMP-1
Specificity:	MMP-1
Other names:	INTERSTITIAL COLLAGENASE
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	3B6
Isotype:	lgG1
Quantity:	0.1 mg

### **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin (1)	-			
ELISA				
Immunoprecipitation			•	
Western Blotting				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein C supernatant	G from tissue culture
Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Ovalbumin conjugated synthetic peptide corresponding to a region within the C-terminus of human MMP-1.
External Database Links	UniProt: P03956 Related reagents  Entrez Gene: 4312 MMP1 Related reagents
Synonyms	CLG
RRID	AB_2144296
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the Ag8563 myeloma cell line.
Specificity	Mouse anti Human MMP-1 antibody, clone 3B6 recognizes human matrix metalloproteinase 1 (MMP-1), also known as interstitial or fibroblast collagenase, a 469 amino acid, in the pro-peptide form, zinc-dependent endopeptidase responsible for degrading the extracellular matrix. MMPs are also involved in cell proliferation, migration, differentiation and apoptosis (Chen et al. 2013). Most MMP's are synthesised as inactive zymogens and a propeptide region must be cleaved off before the enzyme becomes active (Nagase et al. 1992). MMP expression is increased dramatically in a variety of cancer types, where it indicates invasive disease and a poor prognosis (Murray et al. 1996).
Specificity	metalloproteinase 1 (MMP-1), also known as interstitial or fibroblast collagenase, a 469 amino acid, in the pro-peptide form, zinc-dependent endopeptidase responsible for degrading the extracellular matrix. MMPs are also involved in cell proliferation, migration, differentiation and apoptosis (Chen et al. 2013). Most MMP's are synthesised as inactive zymogens and a propeptide region must be cleaved off before the enzyme becomes active (Nagase et al. 1992). MMP expression is increased dramatically in a variety of cancer types, where it indicates invasive disease and a poor prognosis (Murray et al.
Specificity	metalloproteinase 1 (MMP-1), also known as interstitial or fibroblast collagenase, a 469 amino acid, in the pro-peptide form, zinc-dependent endopeptidase responsible for degrading the extracellular matrix. MMPs are also involved in cell proliferation, migration, differentiation and apoptosis (Chen et al. 2013). Most MMP's are synthesised as inactive zymogens and a propeptide region must be cleaved off before the enzyme becomes active (Nagase et al. 1992). MMP expression is increased dramatically in a variety of cancer types, where it indicates invasive disease and a poor prognosis (Murray et al. 1996).  MMP-1 is one of the collagenases, capable of degrading collagens I, II and III, all main components of the interstitial stroma (Robichaud et al. 2011). MMP-1 is overexpressed in invasive melanoma, colorectal and esophageal cancers (Langenskiöld et al. 2013). MMP-1 has also been implicated in arthritis and may influence atherosclerotic lesion formation (Nikkari et al. 1995). Additionally, MMP-1 has been implicated in the repair
Histology Positive Control Tissue	metalloproteinase 1 (MMP-1), also known as interstitial or fibroblast collagenase, a 469 amino acid, in the pro-peptide form, zinc-dependent endopeptidase responsible for degrading the extracellular matrix. MMPs are also involved in cell proliferation, migration, differentiation and apoptosis (Chen et al. 2013). Most MMP's are synthesised as inactive zymogens and a propeptide region must be cleaved off before the enzyme becomes active (Nagase et al. 1992). MMP expression is increased dramatically in a variety of cancer types, where it indicates invasive disease and a poor prognosis (Murray et al. 1996).  MMP-1 is one of the collagenases, capable of degrading collagens I, II and III, all main components of the interstitial stroma (Robichaud et al. 2011). MMP-1 is overexpressed in invasive melanoma, colorectal and esophageal cancers (Langenskiöld et al. 2013). MMP-1 has also been implicated in arthritis and may influence atherosclerotic lesion formation (Nikkari et al. 1995). Additionally, MMP-1 has been implicated in the repair processes of the heart after myocardial infarction (Yarbrough et al. 2003).

References 1. Murray, G.I. et al. (1998) Matrix metalloproteinases and their inhibitors in gastric cancer. Gut. 43: 791-7. 2. Lyall, M.S. et al. (2006) Profiling markers of prognosis in colorectal cancer. Clin Cancer Res. 12: 1184-91. **Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Guarantee 12 months from date of despatch **Health And Safety** Material Safety Datasheet documentation #10040 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA2733GA 10040 Regulatory For research purposes only

### Related Products

## **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (STAR77...)

Rabbit Anti Mouse IgG (STAR12...)

RPE

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) <u>HRP</u>

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

**Recommended Negative Controls** 

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

Email: antibody\_sales\_us@bio-rad.com

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

 ${\bf Email: antibody\_sales\_uk@bio-rad.com}$ 

 ${\bf Email: antibody\_sales\_de@bio-rad.com}$ 

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M405500:220916'

# Printed on 26 Mar 2025