

# Datasheet: MCA2734GA

Description:	MOUSE ANTI HUMAN MMP-3
Specificity:	MMP-3
Other names:	STROMELYSIN-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	1B4
Isotype:	lgG1
Quantity:	0.1 mg

# **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin (1)	•			
ELISA	•			
Immunoprecipitation			•	
Western Blotting				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein supernatant	G from tissue culture
Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Ovalbumin conjugated synthetic peptide corresponding to a region within the C-terminus of human MMP-3.
External Database Links	UniProt: P08254 Related reagents  Entrez Gene: 4314 MMP3 Related reagents
Synonyms	STMY1
RRID	AB_2146579
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the Ag8563 myeloma cell line.
Specificity	Mouse anti Human MMP-3 antibody, clone 1B4 recognizes human matrix metalloproteinase 3, also known as MMP-3, stromelysin-1 or transin-1. MMP-3 in its pro-form is a 477 amino acid ~54kDa zinc dependent endopeptidase processed to a 378 amino acid, ~45 kDa peptide in the mature form containing 4 <a href="https://hemopexin-like">hemopexin-like</a> repeats.  MMPs are also involved in cell proliferation, migration, differentiation and apoptosis (Jana et al. 2012). Most MMP's are synthesised as inactive zymogens and a propeptide region must be cleaved off before the enzyme becomes active. Their expression is increased dramatically in a variety of cancer types, where it indicates invasive disease and a poor prognosis (Holmberg et al. 2013). MMP-3 can degrade proteoglycan, fibronectin, laminin, and type IV collagen, but not interstitial type I collagen (Wilhelm et al. 1993). It can activate other MMP's and may be involved in the repair processes of the heart after myocardial infarction (Wang et al. 2011).  Mouse anti human MMP-3 antibody, clone 1B4 has been used successfully for the identification of MMP-3 in gastric cancer by both Western blotting and
	metalloproteinase 3, also known as MMP-3, stromelysin-1 or transin-1. MMP-3 in its pro-form is a 477 amino acid ~54kDa zinc dependent endopeptidase processed to a 378 amino acid, ~45 kDa peptide in the mature form containing 4 hemopexin-like repeats.  MMPs are also involved in cell proliferation, migration, differentiation and apoptosis (Jana et al. 2012). Most MMP's are synthesised as inactive zymogens and a propeptide region must be cleaved off before the enzyme becomes active. Their expression is increased dramatically in a variety of cancer types, where it indicates invasive disease and a poor prognosis (Holmberg et al. 2013). MMP-3 can degrade proteoglycan, fibronectin, laminin, and type IV collagen, but not interstitial type I collagen (Wilhelm et al. 1993). It can activate other MMP's and may be involved in the repair processes of the heart after myocardial infarction (Wang et al. 2011).  Mouse anti human MMP-3 antibody, clone 1B4 has been used successfully for the identification of MMP-3 in gastric cancer by both Western blotting and immunohistochemistry (Murray et al. 1998)
Histology Positive Control Tissue	metalloproteinase 3, also known as MMP-3, stromelysin-1 or transin-1. MMP-3 in its pro-form is a 477 amino acid ~54kDa zinc dependent endopeptidase processed to a 378 amino acid, ~45 kDa peptide in the mature form containing 4 <a href="https://memory.new.org/new.new.org/">hemopexin-like</a> repeats.  MMPs are also involved in cell proliferation, migration, differentiation and apoptosis ( <a href="Jana et al. 2012">Jana et al. 2012</a> ). Most MMP's are synthesised as inactive zymogens and a propeptide region must be cleaved off before the enzyme becomes active. Their expression is increased dramatically in a variety of cancer types, where it indicates invasive disease and a poor prognosis ( <a href="Holmberg et al. 2013">Holmberg et al. 2013</a> ). MMP-3 can degrade proteoglycan, fibronectin, laminin, and type IV collagen, but not interstitial type I collagen ( <a href="https://wilhelm.et al. 1993">Wilhelm et al. 1993</a> ). It can activate other MMP's and may be involved in the repair processes of the heart after myocardial infarction (

Gut. 43 (6): 791-7.

2. Lyall, M.S. *et al.* (2006) Profiling markers of prognosis in colorectal cancer. <u>Clin Cancer</u> Res. 12 (4): 1184-91.

3. Juica, N.E. *et al.* (2017) *Neisseria gonorrhoeae* Challenge Increases Matrix Metalloproteinase-8 Expression in Fallopian Tube Explants. <u>Front Cell Infect Microbiol. 7:</u> 399.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

 Guarantee
 12 months from date of despatch

 Health And Safety Information
 Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2734GA">https://www.bio-rad-antibodies.com/SDS/MCA2734GA</a>

 Regulatory
 For research purposes only

#### Related Products

## **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (STAR77...)

Rabbit Anti Mouse IgG (STAR12...)

RPE

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Rabbit Anti Mouse IgG (STAR9...) <u>FITC</u>
Goat Anti Mouse IgG (STAR76...) <u>RPE</u>

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

## **Recommended Negative Controls**

#### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M405501:220916'

#### Printed on 12 Aug 2023