# Datasheet: MCA2748F BATCH NUMBER 165480

Description:	RAT ANTI MOUSE CD36:FITC
Specificity:	CD36
Other names:	GPIV (IIIb)
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	MF3
lsotype:	lgG2a
Quantity:	0.1 mg

# **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes No	Not Determined	Suggested Dilution		
	Flow Cytometry	-		Neat		
	necessarily exclude its	use in such proced nmended that the us	ser titrates the product	ng dilutions are given as		
Target Species	Mouse					
Product Form	Purified IgG conjugate	d to Fluorescein Iso	thiocyanate Isomer 1 (F	FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	FITC	490	525			
Preparation	Purified IgG prepared supernatant	by affinity chromato	graphy on Protein G fro	m tissue culture		
Buffer Solution	Phosphate buffered sa	line				
Preservative Stabilisers	0.09% Sodium Azide ( 1% Bovine Serum Albi	6,				
Approx. Protein	IgG concentration 0.1r	ng/ml				

### Concentrations

Immunogen	IL-4 treated murine thioglycollate-elicited peritoneal macrophages
External Database Links	UniProt:         Q08857       Related reagents         Entrez Gene:         12491       Cd36         Related reagents
RRID	AB_1604781
Fusion Partners	Spleen cells from immunised DA rats were fused with cells of the Y3Ag 1.2.3 myeloma cell line
Specificity	<b>Rat anti Mouse CD36 antibody, clone MF3</b> recognizes mouse CD36, also known as platelet glycoprotein 4, glycoprotein IIIb or PAS IV. CD36 is an ~85 kDa multipass transmembrane glycoprotein primarily expressed on platelets, monocytes/macrophages, smooth muscle and endothelial cells. The CD36 molecule is type B scavenger receptor, which binds to multiple ligands including thrombospondin, anionic phospholipids, oxidized low density lipoproteins and long chain fatty acids.
	CD36 has diverse functions and is reported to play a role in innate immunity, platelet adhesion/aggregation and long chain fatty acid transport. The CD36 molecule also directly mediates cytoadhesion of erythrocytes infected with <i>Plasmodium falciparum</i> , and may be involved in the development of atherosclerotic lesions and the formation of foam cells.
	Rat anti Mouse CD36 antibody, clone MF3 has been shown to inhibit IL-4 induced thioglycollate-elicited peritoneal macrophage fusion and significantly block IL-4/GM-CSF-induced bone-marrorw derived macrophage fusion.
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Mwaikambo, B.R. <i>et al.</i> (2009) Hypoxia up-regulates CD36 expression and function via hypoxia-inducible factor-1- and phosphatidylinositol 3-kinase-dependent mechanisms. J Biol Chem. 284: 26695-707.</li> <li>Seeds, R.E. <i>et al.</i> (2011) The role of myeloid receptors on murine plasmacytoid dendritic cells in induction of type I interferon. Int Immunopharmacol. 11: 794-801.</li> <li>Yang, C.N. <i>et al.</i> (2011) Mechanism mediating oligomeric Aβ clearance by naïve primary microglia. Neurobiol Dis. 42 (3): 221-30.</li> <li>Zhou, D. <i>et al.</i> (2012) CD36 level and trafficking are determinants of lipolysis in adipocytes. FASEB J. 26 (11): 4733-42.</li> <li>Samovski, D. <i>et al.</i> (2012) Insulin and AMPK regulate FA translocase/CD36 plasma membrane recruitment in cardiomyocytes via Rab GAP AS160 and Rab8a Rab GTPase. J Lipid Res. 53 (4): 709-17.</li> <li>Dellinger, A. <i>et al.</i> (2013) Functionalization of gadolinium metallofullerenes for detecting atherosclerotic plaque lesions by cardiovascular magnetic resonance. J Cardiovasc Magn</li> </ol>

### Reson. 15: 7.

	7. Miyazaki H et al. (2014) Fatty acid binding protein 7 regulates phagocytosis and			
	cytokine production in Kupffer cells during liver injury. Am J Pathol. 184 (9): 2505-15.			
	8. Pugholm, L.H. et al. (2015) Enhanced Humoral Responses Induced by Targeting of			
	Antigen to Murine Dendritic Cells. Scand J Immunol. 82 (6): 515-22.			
	9. Samovski, D. et al. (2015) Regulation of AMPK activation by CD36 links fatty acid			
	uptake to β-oxidation. Diabetes. 64 (2): 353-9.			
	10. Nie, S. et al. (2015) Detection of atherosclerotic lesions and intimal macrophages			
	using CD36-targeted nanovesicles. <u>J Control Release. pii: S0168-3659(15)30173-5.</u>			
	11. Lund, H. et al. (2018) Competitive repopulation of an empty microglial niche yields			
	functionally distinct subsets of microglia-like cells. Nat Commun. 9 (1): 4845.			
	12. Sui, Y. et al. (2018) Perinatal Bisphenol A Exposure Increases Atherosclerosis in Adult			
	Male PXR-Humanized Mice. Endocrinology. 159 (4): 1595-608.			
	13. Zhang, J. et al. (2019) Anti-atherogenic effects of CD36-targeted epigallocatechin			
	gallate-loaded nanoparticles. <u>J Control Release. 303: 263-73.</u>			
	14. Dhanasekara, C.S. et al. (2021) Nanoparticles target intimal macrophages in			
	atherosclerotic lesions. Nanomedicine. 32: 102346.			
	15. Liu, J. et al. (2022) Pregnane X Receptor Mediates Atherosclerosis Induced by			
	Dicyclohexyl Phthalate in LDL Receptor-Deficient Mice Cells. 11 (7): 1125			
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.			
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in			
	frost-free freezers is not recommended. This product is photosensitive and should be			
	protected from light.			
Guarantee	12 months from date of despatch			
Health And Safety Information	Material Safety Datasheet documentation #10041 available at:			
	https://www.bio-rad-antibodies.com/SDS/MCA2748F			
	10041			
Regulatory	For research purposes only			

## **Related Products**

### **Recommended Negative Controls**

RAT IgG2a NEGATIVE CONTROL:FITC (MCA1212F)

## **Recommended Useful Reagents**

MOUSE SEROBLOCK FcR (BUF041A) MOUSE SEROBLOCK FcR (BUF041B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel
America	Fax: +1 919 878 3751		Fa
	Email: antibody_sales_us@bio-ra	d.com	En

el: +44 (0)1865 852 700 **Europe** ax: +44 (0)1865 852 739 mail: antibody\_sales\_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.com

### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint