

Datasheet: MCA274F

Description:	MOUSE ANTI RAT CD172a:FITC
Specificity:	CD172a
Other names:	SIRP ALPHA
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	OX-41
Isotype:	lgG2a
Quantity:	0.1 mg

## **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat		
Product Form	Purified IgG conjugate	ed to Fluorescein Isoth	hiocyanate Isomer ′
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm
	FITC	490	525
Preparation  Buffer Solution	Purified IgG prepared supernatant  Phosphate buffered s	by affinity chromatogo	raphy on Protein G
Preservative	·	u	
Stabilisers	0.09% Sodium Azide 1.0% Bovine Serum	Albumin	
Approx. Protein			

Immunogen	Rat peritoneal macrophages.	
External Database Links	UniProt:  P97710 Related reagents  Entrez Gene:  25528 Sirpa Related reagents	
Synonyms	Bit, Mfr, Ptpns1, Shps1, Sirp	
RRID	AB_323623	
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with the NS0/U mouse myeloma cell line.	
Specificity	<b>Mouse anti Rat CD172a antibody, clone OX-41</b> recognizes rat CD172a, also known as Signal regulatory protein (SIRP). CD172a is expressed selectively by myeloid cells and by neurons.	
	MRC OX-41 binds to a different epitope on SIRP than ED9. Mouse anti Rat CD172a antibody, clone OX-41 does not completely block the interaction of CD172a - CD47 (Vernon-Wilson et al. 2000).	
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.	
References	<ol> <li>Robinson, A.P. <i>et al.</i> (1986) Macrophage heterogeneity in the rat as delineated by two monoclonal antibodies MRC OX-41 and MRC OX-42, the latter recognizing complement receptor type 3. Immunology. 57 (2): 239-47.</li> <li>Santos, L.L. <i>et al.</i> (1997) Anti-neutrophil monoclonal antibody therapy inhibits the development of adjuvant arthritis. Clin Exp Immunol. 107: 248-53.</li> <li>Adams, S. <i>et al.</i> (1998) Signal-regulatory protein is selectively expressed by myeloid and neuronal cells. J Immunol. 161 (4): 1853-9.</li> <li>Milling, S.W. <i>et al.</i> (2006) Collection of lymph-borne dendritic cells in the rat. Nat Protoc. 1: 2263-70.</li> <li>Moghaddami, M. <i>et al.</i> (2007) Recruitment of dendritic cells and macrophages during T cell-mediated synovial inflammation. Arthritis Res Ther. 9: R120.</li> <li>Varalakshmi, Ch, <i>et al.</i> (2008) Immunomodulatory effects of curcumin: <i>in-vivo</i>. Int Immunopharmacol. 8 (5): 688-700.</li> <li>Zhou, Y.J. <i>et al.</i> (2010) The role of the lactadherin in promoting intestinal DCs development <i>in vivo</i> and <i>invitro</i>. Clin Dev Immunol. 2010: 357541.</li> </ol>	
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.	
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be	

## protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA274F">https://www.bio-rad-antibodies.com/SDS/MCA274F</a> 10041
Regulatory	For research purposes only

# **Related Products**

# **Recommended Negative Controls**

MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA1210F)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385558:210513'

## Printed on 27 Aug 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint