

## Datasheet: MCA274G

#### **BATCH NUMBER 1710**

Description: MOUSE ANTI RAT CD172		
Specificity:	CD172a	
Other names:	SIRP ALPHA	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	OX-41	
Isotype:	IgG2a	
Quantity:	1 mg	

## **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/100 - 1/200
Immunohistology - Frozen	•			
Immunohistology - Paraffin (1)	•			
ELISA			•	
Immunoprecipitation			•	
Western Blotting				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)This product requires protein digestion pre-treatment of paraffin sections e.g. trypsin or pronase.

Target Species	Rat	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein G supernatant	G from tissue culture
Buffer Solution	Phosphate buffered saline	

0.09% Sodium Azide
Yes
IgG concentration 1.0 mg/ml
Rat peritoneal macrophages.
UniProt: P97710 Related reagents  Entrez Gene: 25528 Sirpa Related reagents
Bit, Mfr, Ptpns1, Shps1, Sirp
AB_323480
Spleen cells from immunised BALB/c mice were fused with the NS0/U mouse myeloma cell line.
<b>Mouse anti Rat CD172a antibody, clone OX-41</b> recognizes rat CD172a, also known as Signal regulatory protein (SIRP). CD172a is expressed selectively by myeloid cells and by neurons.
MRC OX-41 binds to a different epitope on SIRP than ED9. It does not block the interaction of CD172a - CD47.
Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<ol> <li>Robinson, A.P. <i>et al.</i> (1986) Macrophage heterogeneity in the rat as delineated by two monoclonal antibodies MRC OX-41 and MRC OX-42, the latter recognizing complement receptor type 3. <a href="Immunology.57">Immunology.57</a> (2): 239-47.</li> <li>Barclay, A.N. (1981) The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. <a href="Immunology.42">Immunology.42</a> (4): 593-600.</li> <li>Adams, S. <i>et al.</i> (1998) Signal-regulatory protein is selectively expressed by myeloid and neuronal cells. <a href="Jenance-Jimmunol.161">Jimmunol.161</a> (4): 1853-9.</li> <li>Santos, L.L. <i>et al.</i> (1997) Anti-neutrophil monoclonal antibody therapy inhibits the development of adjuvant arthritis. <a href="Clin Exp Immunol.107">Clin Exp Immunol. 107</a>: 248-53.</li> <li>Milling, S.W. <i>et al.</i> (2006) Collection of lymph-borne dendritic cells in the rat. <a href="Nat Protocoli:2263-70">Nat Protocoli:2263-70</a>.</li> <li>Moghaddami, M. <i>et al.</i> (2007) Recruitment of dendritic cells and macrophages during T cell-mediated synovial inflammation. <a href="Arthritis Res Ther.9">Arthritis Res Ther.9</a>: R120.</li> <li>Zhou, Y.J. <i>et al.</i> (2010) The role of the lactadherin in promoting intestinal DCs development <i>in vivo</i> and <i>invitro</i>. <a href="Clin Dev Immunol.2010:357541">Clin Dev Immunol.2010:357541</a>.</li> </ol>

8. Varalakshmi, Ch, et al. (2008) Immunomodulatory effects of curcumin: in-vivo. Int

### Immunopharmacol. 8 (5): 688-700.

Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted.
	Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA274G">https://www.bio-rad-antibodies.com/SDS/MCA274G</a> 10040
Regulatory	For research purposes only

### **Related Products**

# **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...) **RPE** Goat Anti Mouse IgG IgA IgM (STAR87...) HRP Goat Anti Mouse IgG (STAR76...) **RPE** Rabbit Anti Mouse IgG (STAR13...) **HRP** 

Goat Anti Mouse IgG (STAR70...) **FITC** 

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) **FITC** Goat Anti Mouse IgG (STAR77...) **HRP** 

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

## **Recommended Negative Controls**

America

## MOUSE IgG2a NEGATIVE CONTROL (MCA1210)

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700

> Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 Fax: +1 919 878 3751

Email: antibody\_sales\_us@bio-rad.com Email: antibody\_sales\_uk@bio-rad.com Email: antibody\_sales\_de@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M367412:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint