

Datasheet: MCA275R

Description:	MOUSE ANTI RAT CD11b
Specificity:	CD11b
Other names:	INTEGRIN ALPHA M CHAIN, MAC-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	OX-42
Isotype:	IgG2a
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/50 - 1/100
Immunohistology - Paraffin (1)			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative /positive controls.

(1)**OX-42 is reported to be suitable for paraffin-embedded sections following PLP fixation ([Whiteland et al., 1995](#)).**

Target Species	Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Resident rat peritoneal macrophages.
RRID	AB_321302
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the NSO/U mouse myeloma cell line.
Specificity	<p>Mouse anti Rat CD11b, clone OX-42, recognizes rat CD11b, also known as integrin alpha-M, the receptor for the iC3b component of complement. CD11b is a 1151 amino acid single pass type 1 transmembrane glycoprotein possessing a single vWFA domain and multiple FG-GAP repeats. CD11b is expressed on most macrophages, including resident and activated peritoneal macrophages and Kupffer cells and around 35% of alveolar macrophages. The antibody also labels dendritic cells, granulocytes and microglia in the brain (Robinson et al.1986).</p> <p>Mouse anti Rat CD11b, clone OX-42 is reported to inhibit complement mediated rosettes (Robinson et al.1986) as well as inhibit myelin binding and uptake (van der Laan et al.1996).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Robinson, A.P. <i>et al.</i> (1986) Macrophage heterogeneity in the rat as delineated by two monoclonal antibodies MRC OX-41 and MRC OX-42, the latter recognizing complement receptor type 3. Immunology. 57 (2): 239-47. 2. Barclay, A.N. (1981) The localization of population of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. Immunology. 42: 593-600. 3. Milligan, C.E. <i>et al.</i> (1991) Differential immunochemical markers reveal the normal distribution of brain macrophages and microglia in the developing rat brain. J Comp Neurol. 314 (1): 125-35. 4. Whiteland, J.L. <i>et al.</i> (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem. 43 (3): 313-20. 5. Zilka, N. <i>et al.</i> (2009) Human misfolded truncated tau protein promotes activation of microglia and leukocyte infiltration in the transgenic rat model of tauopathy. J Neuroimmunol. 209 (1-2): 16-25. 6. Shokouhi, B.N. <i>et al</i> (2010) Microglial responses around intrinsic CNS neurons are correlated with axonal regeneration. BMC Neurosci. 11: 13. 7. Calvo, M. <i>et al.</i> (2010) Neuregulin-ErbB signaling promotes microglial proliferation and chemotaxis contributing to microgliosis and pain after peripheral nerve injury. J Neurosci. 30 (15): 5437-50.

8. Jean, Y.H. *et al.* (2009) Capnellene, a natural marine compound derived from soft coral, attenuates chronic constriction injury-induced neuropathic pain in rats [Br J Pharmacol. 158: 713-25.](#)
9. Morales-Garcia, J.A. *et al.* (2011) Phosphodiesterase 7 inhibition preserves dopaminergic neurons in cellular and rodent models of Parkinson disease. [PLoS One. 6\(2\):e17240.](#)
10. Spencer-Segal, J.L. *et al.* (2011) Distribution of Phosphorylated TrkB Receptor in the Mouse Hippocampal Formation Depends on Sex and Estrous Cycle Stage. [J Neurosci. 31: 6780-90.](#)
11. Chew, S.S. *et al.* (2011) Response of retinal Connexin43 to optic nerve injury. [Invest Ophthalmol Vis Sci. 52: 3620-9.](#)
12. Huh, S.H. *et al.* (2011) Ethyl pyruvate rescues nigrostriatal dopaminergic neurons by regulating glial activation in a mouse model of Parkinson's disease. [J Immunol. 187: 960-9.](#)
13. Jeong, H.K. *et al.* (2010) Inflammatory responses are not sufficient to cause delayed neuronal death in ATP-induced acute brain injury. [PLoS One. 5: e13756.](#)
14. Liew, H.K. *et al.* (2012) Systemic administration of urocortin after intracerebral hemorrhage reduces neurological deficits and neuroinflammation in rats. [J Neuroinflammation. 9: 13.](#)
15. Szmydynger-Chodobska, J. *et al.* (2011) Multiple sites of vasopressin synthesis in the injured brain. [J Cereb Blood Flow Metab. 31: 47-51.](#)
16. Tchoukalova, Y.D. *et al.* (2012) *In vivo* adipogenesis in rats measured by cell kinetics in adipocytes and plastic-adherent stroma-vascular cells in response to high-fat diet and thiazolidinedione. [Diabetes. 61: 137-44.](#)
17. Xu, Q. *et al.* (2011) Spinal phosphoinositide 3-kinase-Akt-mammalian target of rapamycin signaling cascades in inflammation-induced hyperalgesia. [J Neurosci. 31: 2113-24.](#)
18. Lovett-Barr, M.R. *et al.* (2012) Repetitive intermittent hypoxia induces respiratory and somatic motor recovery after chronic cervical spinal injury. [J Neurosci. 32 \(11\): 3591-600.](#)
19. Lavis, S. *et al.* (2012) Reactive astrocytes overexpress TSPO and are detected by TSPO positron emission tomography imaging. [J Neurosci. 32: 10809-18.](#)
20. Ortega, F.J. *et al.* (2012) Glibenclamide enhances neurogenesis and improves long-term functional recovery after transient focal cerebral ischemia. [J Cereb Blood Flow Metab. 33: 356-64.](#)
21. Zhao, H. *et al.* (2013) Brain 3-Mercaptopyruvate Sulfurtransferase (3MST): Cellular Localization and Downregulation after Acute Stroke. [PLoS One. 8\(6\):e67322.](#)
22. Huang S *et al.* (2015) Expression of Peroxiredoxin 1 After Traumatic Spinal Cord Injury in Rats. [Cell Mol Neurobiol. May 24. \[Epub ahead of print\]](#)
23. Wang, A.Y. *et al.* (2015) Searching for a rat model of chronic tympanic membrane perforation: Healing delayed by mitomycin C/dexamethasone but not paper implantation or iterative myringotomy. [Int J Pediatr Otorhinolaryngol. 79 \(8\): 1240-7.](#)
24. Hernangómez M *et al.* (2016) CD200R1 agonist attenuates glial activation, inflammatory reactions, and hypersensitivity immediately after its intrathecal application in a rat neuropathic pain model. [J Neuroinflammation. 13 \(1\): 43.](#)
25. Signarovitz, A.L. *et al.* (2012) Mucosal immunization with live attenuated *Francisella novicida* U112ΔiglB protects against pulmonary *F. tularensis* SCHU S4 in the Fischer 344 rat model. [PLoS One. 7: e47639.](#)

26. Liu, Z. *et al.* (2016) Leukocyte Infiltration Triggers Seizure Recurrence in a Rat Model of Temporal Lobe Epilepsy. [Inflammation. Apr 4. \[Epub ahead of print\].](#)
27. Szmydynger-Chodobska, J. *et al.* (2016) The Involvement of Pial Microvessels in Leukocyte Invasion after Mild Traumatic Brain Injury. [PLoS One. 11 \(12\): e0167677.](#)
28. Ejaz, S. *et al.* (2014) MRI and neuropathological validations of the involvement of air pollutants in cortical selective neuronal loss. [Environ Sci Pollut Res Int. 21 \(5\): 3351-62.](#)
29. Alizadeh, A. *et al.* (2017) Neuregulin-1 positively modulates glial response and improves neurological recovery following traumatic spinal cord injury. [Glia. Apr 29. \[Epub ahead of print\]](#)
30. Popiolek-Barczyk, K. *et al.* (2017) Biphalin, a Dimeric Enkephalin, Alleviates LPS-Induced Activation in Rat Primary Microglial Cultures in Opioid Receptor-Dependent and Receptor-Independent Manners. [Neural Plasticity. 2017: 1-19.](#)
31. Huang RY *et al.* (2017) Rapid and Delayed Effects of Pulsed Radiofrequency on Neuropathic Pain: Electrophysiological, Molecular, and Behavioral Evidence Supporting Long-Term Depression. [Pain Physician. 20 \(2\): E269-E283.](#)
32. Bourke, G. *et al.* (2017) Effects of early nerve repair on experimental brachial plexus injury in neonatal rats. [J Hand Surg \(Eur Vol\). 1753193417732696 \[Epub ahead of print\].](#)
33. Wang, Z.C. *et al.* (2017) Involvement of NF- κ B and the CX3CR1 Signaling Network in Mechanical Allodynia Induced by Tetanic Sciatic Stimulation. [Neurosci Bull. Jun 13 \[Epub ahead of print\].](#)
34. Klemm, P. *et al.* (2019) Hypothermia protects retinal ganglion cells against hypoxia-induced cell death in a retina organ culture model. [Clin Exp Ophthalmol. Jun 01 \[Epub ahead of print\].](#)
35. de Sousa, É *et al.* (2013) Developmental and functional expression of miRNA-stability related genes in the nervous system. [PLoS One. 8 \(5\): e56908.](#)
36. van Vliet, E.A. (2020) Long-lasting blood-brain barrier dysfunction and neuroinflammation after traumatic brain injury [Neurobiol Dis 11:13.](#)
37. Kuo, T. *et al.* (2020) Post-stroke Delivery of Valproic acid Promotes Functional Recovery and Differentially Modifies Responses of Peri-infarct Microglia. [Res Sq; Dec 01. \[Preprint Epub ahead of print\].](#)
38. Szeredi, I.D. *et al.* (2020) Prior perineural or neonatal treatment with capsaicin does not alter the development of spinal microgliosis induced by peripheral nerve injury. [Cell Tissue Res. Sep 22 \[Epub ahead of print\].](#)
39. Lin, J. *et al.* (2020) Protective effect of Soluble Epoxide Hydrolase Inhibition in Retinal Vasculopathy associated with Polycystic Kidney Disease. [Theranostics. 10 \(17\): 7857-71.](#)
40. Tanaka, J. *et al.* (2020) Generation of CSF1-Independent Ramified Microglia-Like Cells from Leptomeninges *In Vitro*.. [Cells. 10 \(1\) Dec 25 \[Epub ahead of print\].](#)
41. Espinosa-Garcia, C. *et al.* (2020) Progesterone Attenuates Stress-Induced NLRP3 Inflammasome Activation and Enhances Autophagy following Ischemic Brain Injury. [Int J Mol Sci. 21 \(11\): 3740.](#)
42. Memedovski, Z. *et al.* (2020) Classical and Alternative Activation of Rat Microglia Treated with Ultrapure *Porphyromonas gingivalis* Lipopolysaccharide *In Vitro*.. [Toxins \(Basel\). 12\(5\): 333.](#)
43. Mecha, M. *et al.* (2020) Involvement of Wnt7a in the role of M2c microglia in neural stem cell oligodendrogenesis. [J Neuroinflammation. 17 \(1\): 88.](#)
44. Terayama, R. *et al.* (2018) A₃ adenosine receptor agonist attenuates neuropathic pain by suppressing activation of microglia and convergence of nociceptive inputs in the spinal

- dorsal horn. [Exp Brain Res. 236 \(12\): 3203-13.](#)
45. Sugama, S. *et al.* (2019) Stress-induced microglial activation occurs through β -adrenergic receptor: noradrenaline as a key neurotransmitter in microglial activation. [J Neuroinflammation. 16 \(1\): 266.](#)
46. Winkler, A. *et al.* (2021) Blood-brain barrier resealing in neuromyelitis optica occurs independently of astrocyte regeneration. [J Clin Invest. 131\(5\):e141694.](#)
47. Hosoi, R. *et al.* (2021) Evaluation of Intracellular Processes in Quinolinic Acid-Induced Brain Damage by Imaging Reactive Oxygen Species Generation and Mitochondrial Complex I Activity. [Res Square.10 May \[Epub ahead of print\].](#)
48. Hellenbrand, D.J. *et al.* (2019) Sustained interleukin-10 delivery reduces inflammation and improves motor function after spinal cord injury. [J Neuroinflammation. 16 \(1\): 93.](#)
49. Collins, J.J.P. *et al.* (2018) Impaired Angiogenic Supportive Capacity and Altered Gene Expression Profile of Resident CD146⁺ Mesenchymal Stromal Cells Isolated from Hypoxia-Injured Neonatal Rat Lungs. [Stem Cells Dev. 27 \(16\): 1109-24.](#)
50. Sanches, E.F. *et al.* (2018) Brain Metabolism Alterations Induced by Pregnancy Swimming Decreases Neurological Impairments Following Neonatal Hypoxia-Ischemia in Very Immature Rats. [Front Neurol. 9: 480.](#)
51. Hahm, S.C. *et al.* (2019) Transcutaneous Electrical Nerve Stimulation Reduces Knee Osteoarthritic Pain by Inhibiting Spinal Glial Cells in Rats. [Phys Ther. 99 \(9\): 1211-23.](#)
52. Muratori, L. *et al.* (2019) New basic insights on the potential of a chitosan-based medical device for improving functional recovery after radical prostatectomy. [BJU Int. 124 \(6\): 1063-76.](#)
53. Giuliano, C. *et al.* (2021) Neuroprotective and Symptomatic Effects of Cannabidiol in an Animal Model of Parkinson's Disease. [Int J Mol Sci. 22 \(16\): 8920.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Goat Anti Mouse IgG (STAR76...)	RPE
Human Anti Mouse IgG2a (HCA037...)	HRP

Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®550](#),
[DyLight®650](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA1210\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M383970:210513'

Printed on 10 Sep 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)