

Datasheet: MCA2804C

BATCH NUMBER 155679

Description:	MOUSE ANTI HUMAN CD14:RPE-Cy5
Specificity:	CD14
Format:	RPE-CY5
Product Type:	Monoclonal Antibody
Clone:	61D3
Isotype:	IgG1
Quantity:	100 TESTS/1ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Bovine, Dog, Rabbit, Sheep, Pig, Cynomolgus monkey, Goat, Cat, Mink
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - Cy5 - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE-Cy5 488nm laser	496	667

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.1% Sodium Azide (NaN₃)
 Sucrose

External Database
Links

UniProt:

[P08571](#) [Related reagents](#)

Entrez Gene:

[929](#) CD14 [Related reagents](#)

RRID

AB_1100736

Specificity

Mouse anti Human CD14 antibody, clone 61D3 recognizes human CD14, otherwise known as monocyte differentiation antigen. It is a ~40 kDa protein found on cell surfaces, particularly macrophages. CD14 acts as a co-receptor (along with the Toll-like receptor TLR 4 and MD-2) to mediate the innate immune response to bacterial lipopolysaccharide.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells or 100ul whole blood.

References

1. Gagro, A. *et al.* (2006) Type I cytokine profiles of human naïve and memory B lymphocytes: a potential for memory cells to impact polarization. [Immunology. 118 \(1\): 66-77.](#)
2. Barat, C. *et al.* (2008) Extracellular ATP reduces HIV-1 transfer from immature dendritic cells to CD4+ T lymphocytes. [Retrovirology. 5: 30.](#)
3. Levy, O. *et al.* (2003) Critical role of the complement system in group B *streptococcus*-induced tumor necrosis factor alpha release. [Infect Immun. 71 \(11\): 6344-53.](#)
4. Raghuraman, S. *et al.* (2012) Spontaneous clearance of chronic hepatitis C virus infection is associated with appearance of neutralizing antibodies and reversal of T-cell exhaustion. [J Infect Dis. 205: 763-71.](#)
5. Balasubramanian, K. & Schroit, A.J. (1998) Characterization of phosphatidylserine-dependent beta2-glycoprotein I macrophage interactions. Implications for apoptotic cell clearance by phagocytes. [J Biol Chem. 273 \(44\): 29272-7.](#)
6. Eleftheriou D *et al.* (2012) Endothelial injury in childhood stroke with cerebral arteriopathy: a cross-sectional study. [Neurology. 79 \(21\): 2089-96.](#)
7. Henriksen, P.A. *et al.* (2004) Gene delivery of the elastase inhibitor elafin protects macrophages from neutrophil elastase-mediated impairment of apoptotic cell recognition. [FEBS Lett. 574: 80-4.](#)
8. Giles, K.M. *et al.* (2001) Glucocorticoid augmentation of macrophage capacity for phagocytosis of apoptotic cells is associated with reduced p130Cas expression, loss of paxillin/pyk2 phosphorylation, and high levels of active Rac. [J Immunol. 167: 976-86.](#)
9. Amorim, I.F. *et al.* (2011) Toll receptors type-2 and CR3 expression of canine monocytes and its correlation with immunohistochemistry and xenodiagnosis in visceral leishmaniasis. [PLoS One. 6: e27679.](#)
10. Werner, J.M. *et al.* (2013) Innate immune responses in hepatitis C virus-exposed healthcare workers who do not develop acute infection. [Hepatology. 58 \(5\): 1621-31.](#)
11. Santos, B.P. *et al.* (2017) Blood and milk polymorphonuclear leukocyte and monocyte/macrophage functions in naturally caprine arthritis encephalitis virus infection in dairy goats. [Vet Immunol Immunopathol. 188: 21-6.](#)
12. Serti, E. *et al.* (2015) Successful Interferon-Free Therapy of Chronic Hepatitis C Virus Infection Normalizes Natural Killer Cell Function. [Gastroenterology. 149 \(1\): 190-200.e2.](#)

Storage Store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee Guaranteed until date of expiry. Please see product label.

Acknowledgements Cy® and CyDye® are registered trademarks of GE Healthcare

Health And Safety Information Material Safety Datasheet documentation #10331 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA2804C>
10331

Regulatory For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M350421:190307'

Printed on 25 Mar 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)