

Datasheet: MCA2806SBV760

| Description: | MOUSE ANTI HUMAN CD69:StarBright Violet 760 | | |
|---------------|---|--|--|
| Specificity: | CD69 | | |
| Other names: | AIM | | |
| Format: | StarBright Violet 760 | | |
| Product Type: | Monoclonal Antibody | | |
| Clone: | FN50 | | |
| Isotype: | IgG1 | | |
| Quantity: | 100 TESTS/0.5ml | | |
| | | | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | • | | | Neat |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| Target Species | Human | | | |
|-----------------------------|---|---|--|---|
| Species Cross Reactivity | N.B. Antibody reactive reactivity is derived for | rity and working condition testing within our I | ions may vary betwaboratories, peer-re | esus Monkey, Macaque veen species. Cross eviewed publications or references indicated fo |
| Product Form | Purified IgG conjugat | ted to StarBright Violet | 760 - liquid | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nn | n) |
| | StarBright Violet 760 | 403 | 754 | |
| Preparation | Purified IgG prepared supernatant | d by affinity chromatog | raphy on Protein G | from tissue culture |
| Buffer Solution | Phosphate buffered s | saline | | |

Preservative Stabilisers

0.09% Sodium Azide (NaN₃)1% Bovine Serum Albumin

0.1% Pluronic F680.1% PEG 33500.05% Tween 20

Approx. Protein Concentrations

For information on the concentration of our StarBright Dye conjugated reagents please visit our FAQ page.

Immunogen

Activated human B-cells.

External Database Links

UniProt:

Q07108 Related reagents

Entrez Gene:

969 CD69 Related reagents

Synonyms

CLEC2C

Specificity

Mouse anti Human CD69 antibody, clone FN50 recognizes the human early activation antigen CD69, also known as activation inducer molecule (AIM), Early T-cell activation antigen p60, EA1 or MLR-3. CD69 is a 199 amino acid single pass type II transmembrane glycoprotein of ~30 kDa containing a single C-type lectin domain and a single potential N-glycosylation site. CD69 is expressed as a disulphide bond linked homodimer of ~60 kDa (López-Cabrera et al. 1993).

CD69 is a marker of early activation expressed by B and T lymphocytes, natural killer cells(<u>Werfel 1997</u>), neutrophils, thymocytes and platelets (<u>Gaviol et al. 1992</u>). Expression of CD69 is rapidly induced on activation by infection or chronic inflamation (<u>Sancho et al. 2005</u>). Multiple dimeric glycoforms of CD69 can be formed through differential glycosylation of the monomeric subunits (<u>Vance et al. 1997</u>).

Mouse anti Human CD69 , clone FN50 is useful for the detection of CD69 by flow cytometry and immunohistochemistry on frozen tissue sections.

Flow Cytometry

Use 5ul of the suggested working dilution to label 10⁶ cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Holte, H. *et al.* (1989) Ki67 and 4F2 antigen expression as well as DNA synthesis predict survival at relapse/tumour progression in low-grade B-cell lymphoma. <u>Int J Cancer.</u> 44 (6): 975-80.
- 2. Herberth, M. *et al.* (2010) Differential effects on T-cell function following exposure to serum from schizophrenia smokers. Mol Psychiatry. 15 (4): 364-71.
- 3. Schaeuble, K. *et al.* (2011) Cross-talk between TCR and CCR7 signaling sets a temporal threshold for enhanced T lymphocyte migration. J Immunol. 187 (11): 5645-52.
- 4. Sela, M. *et al.* (2011) Sequential phosphorylation of SLP-76 at tyrosine 173 is required for activation of T and mast cells. <u>EMBO J. 30 (15): 3160-72.</u>

- 5. Garbe, Y. et al. (2011) Semiallogenic fusions of MSI(+) tumor cells and activated B cells induce MSI-specific T cell responses. BMC Cancer. 11: 410.
- 6. Schwitalle, Y. et al. (2004) Immunogenic peptides generated by frameshift mutations in DNA mismatch repair-deficient cancer cells. Cancer Immun. 4: 14.
- 7. Sutavani, R.V. et al. (2013) CD55 Costimulation Induces Differentiation of a Discrete T Regulatory Type 1 Cell Population with a Stable Phenotype. J Immunol. 191: 5895-903.
- 8. Walter, G.J. et al. (2013) Interaction with activated monocytes enhances cytokine expression and suppressive activity of human CD4+CD45ro+CD25+CD127(low) regulatory T cells. Arthritis Rheum. 65: 627-38.
- 9. Kuric, E. et al. (2017) Demonstration of Tissue Resident Memory CD8 T Cells in Insulitic Lesions in Adult Patients with Recent-Onset Type 1 Diabetes. Am J Pathol. 187 (3): 581-8.
- 10. Karnell, F.G. et al. (2017) Reconstitution of immune cell populations in multiple sclerosis patients after autologous stem cell transplantation. Clin Exp Immunol. 189 (3): 268-278.
- 11. Rossatti, P. et al. (2022) Rapid increase in transferrin receptor recycling promotes adhesion during T cell activation. BMC Biol. 20 (1): 189.

| Storage | Store at +4°C. DO NOT FREEZE. |
|-------------------|---|
| | This product should be stored undiluted. |
| Guarantee | 12 months from date of despatch |
| Acknowledgements | This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts |
| Health And Safety | Material Safety Datasheet documentation #20471 available at: |
| Information | https://www.bio-rad-antibodies.com/SDS/MCA2806SBV760 |
| | 20471 |
| Regulatory | For research purposes only |

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376 America

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Email: antibody sales de@bio-rad.com

Email: antibody sales us@bio-rad.com

Email: antibody sales uk@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M437328:250311'

Printed on 03 Apr 2025