

## Datasheet: MCA2808C

<b>Description:</b>	MOUSE ANTI HUMAN CD25:RPE-Cy5
<b>Specificity:</b>	CD25
<b>Other names:</b>	IL-2R ALPHA CHAIN
<b>Format:</b>	RPE-CY5
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	BC96
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	100 TESTS/0.5ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

### Target Species

Human

### Species Cross Reactivity

Reacts with: Cynomolgus monkey, Rhesus Monkey

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

### Product Form

Purified IgG conjugated to RPE-Cy5 - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE-Cy5 488nm laser	496	667

### Preparation

Purified IgG prepared by affinity chromatography

### Buffer Solution

Phosphate buffered saline

<b>Preservative</b>	0.09% Sodium Azide
<b>Stabilisers</b>	0.2% Bovine Serum Albumin
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P01589</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3559</a>    IL2RA    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_1101641
<b>Specificity</b>	<p><b>Mouse anti Human CD25 antibody, clone BC96</b> recognizes the human CD25 cell surface antigen, also known as the Interleukin-2 receptor subunit alpha or TAC antigen.</p> <p>CD25 is derived from a 272 amino acid precursor to yield a 251 amino acid monomeric receptor of ~53kDa which acts as the low affinity receptor for IL-2. CD25 can also form a heterodimer with <a href="#">CD122</a> to form the high affinity receptor for IL-2. CD25 is expressed by activated lymphocytes (<a href="#">Jackson et al. 1990</a>, <a href="#">Butcher et al. 1990</a>). Defects in CD4+ CD25+ regulatory T cells have been implicated in the pathogenesis of autoimmune diabetes (<a href="#">Lindley et al. 2005</a>) and other autoimmune conditions (<a href="#">Sakaguchi et al. 1995</a>).</p>
<b>Flow Cytometry</b>	Use 5ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Chapel, A. <i>et al.</i> (1992) Differential human immunodeficiency virus expression in CD4+ cloned lymphocytes: from viral latency to replication. <a href="#">J Virol. 66 (6): 3966-70.</a></li> <li>2. Barberá A <i>et al.</i> (2013) APL-1, an altered peptide ligand derived from human heat-shock protein 60, selectively induces apoptosis in activated CD4+ CD25+ T cells from peripheral blood of rheumatoid arthritis patients. <a href="#">Int Immunopharmacol. 17 (4): 1075-83.</a></li> <li>3. Duggleby, R.C. &amp; Madrigal, J.A. (2014) Methods of detection of immune reconstitution and T regulatory cells by flow cytometry. <a href="#">Methods Mol Biol. 1109: 159-86.</a></li> </ol>
<b>Storage</b>	<p>Store at +4°C.</p> <p>DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
<b>Guarantee</b>	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.
<b>Acknowledgements</b>	Cy® and CyDye® are registered trademarks of GE Healthcare
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA2808C">https://www.bio-rad-antibodies.com/SDS/MCA2808C</a>

---

**Regulatory**For research purposes only

---

**Related Products****Recommended Useful Reagents**[HUMAN SEROBLOCK \(BUF070A\)](#)[HUMAN SEROBLOCK \(BUF070B\)](#)**North & South** Tel: +1 800 265 7376**America** Fax: +1 919 878 3751Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M361253:200210'

**Printed on 18 Jan 2024**