

Datasheet: MCA2814C

Description:	MOUSE ANTI HUMAN CD2:RPE-Cy5
Specificity:	CD2
Other names:	LFA-2
Format:	RPE-CY5
Product Type:	Monoclonal Antibody
Clone:	RPA-2.10
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Species Cross Reactivity	Reacts with: Macaque, Cynomolgus monkey, Rhesus Monkey, Pig, Baboon, Chimpanzee N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to RPE-Cy5 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE-Cy5 488nm laser	496	667
Preparation	Purified IgG prepared by affinity chromatography		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	0.2% Bovine Serum Albumin		

External Database Links

UniProt:

[P06729](#) [Related reagents](#)

Entrez Gene:

[914](#) CD2 [Related reagents](#)

Synonyms	SRBC
RRID	AB_1101170
Specificity	Mouse anti Human CD2 antibody, clone RPA-2.10 recognizes the human CD2 cell surface antigen. CD2 is expressed by T lymphocytes and by NK cells.
Flow Cytometry	Use 5ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood.
References	<ol style="list-style-type: none"> 1. Scherthner, G.H. <i>et al.</i> (2001) Expression, epitope analysis, and functional role of the LFA-2 antigen detectable on neoplastic mast cells. Blood. 98 (13): 3784-92. 2. Aversa GG <i>et al.</i> (1987) RPA-2.10: an anti-CD2 monoclonal antibody that inhibits alloimmune responses and monitors T cell activation. Transplant Proc. 19 (1 Pt 1): 277-8.
Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54.
Storage	<p>Store at +4°C.</p> <p>DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.
Acknowledgements	Cy® and CyDye® are registered trademarks of GE Healthcare
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M350509:190307'

Printed on 11 Aug 2020