

# Datasheet: MCA2873

Description: MOUSE ANTI RAT CD		
Specificity:	CD80	
Other names:	B7-1	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	3H5	
lsotype:	lgG1	
Quantity:	0.25 mg	

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is				
	derived from testing within our laboratories, peer-reviewed publications or personal				
	communications from the originators. Please refer to references indicated for further				
	information. For general protocol recommendations, please visit <u>www.bio-</u>				
	rad-antibodies.com/protocols.				

		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	-			1/25 - 1/200
	Immunohistology - Frozen	-			
	Immunohistology - Paraffin			•	
	ELISA				
	Immunoprecipitation	-			
	Western Blotting				
	Where this product has n	ot been te	ested for	use in a particular tec	hnique this does not
	necessarily exclude its use in such procedures. Suggested working dilutions are given a				
	a guide only. It is recommended that the user titrates the product for use in their own				
	system using appropriate			•	
Target Species	Rat				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant			raphy on Protein A fro	m tissue culture
Buffer Solution	Phosphate buffered salin	е			
Preservative Stabilisers	0.09% Sodium Azide (Na	ιN <sub>3</sub> )			

Carrier Free	Yes			
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml			
Immunogen	HTLV-1 transformed Lewis-S1 cells.			
External Database Links	UniProt: <u>O55202</u> <u>Related reagents</u>			
RRID	AB_1720046			
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the P3U1 mouse myeloma cell line.			
Specificity	<ul> <li>Mouse anti Rat CD80, clone 3H5 specifically recognizes rat CD80, otherwise known as B7-1, a type I transmembrane glycoprotein and member of the Ig superfamily, which acts as a ligand for both CD28 and CD152 (CTLA-4), and is primarily expressed on antigen presenting cells (APCs) including dendritic cells.</li> <li>CD80 is a B cell activation antigen, which functions in the CD28-CD80/CD86 co-stimulatory pathway vital for T cell activation and proliferation. In contrast, the interaction of CD80 with CD152 has an inhibitory effect on T cell responses.</li> </ul>			
	Clone 3H5 has been shown to block the co-stimulatory activity of rat CD80.			
Flow Cytometry	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.			
References	<ol> <li>Maeda, K. <i>et al.</i> (1997) Characterization of rat CD80 and CD86 by molecular cloning and mAb. <u>Int. Immunol. 9: 993-1000.</u></li> <li>Damoiseaux, J.G. <i>et al.</i> (1998) Costimulatory molecules CD80 and CD86 in the rat; tissue distribution and expression by antigen-presenting cells. <u>J Leukoc Biol. 64 (6)</u>: <u>803-9.</u></li> <li>Kano, M. <i>et al.</i> (1998) A crucial role of host CD80 and CD86 in rat cardiac xenograft rejection in mice. <u>Transplantation. 65: 837-43.</u></li> <li>Hanabuchi, S. <i>et al.</i> (2000) Development of human T-cell leukemia virus type 1-transformed tumors in rats following suppression of T-cell immunity by CD80 and CD86 blockade. <u>J Virol. 74: 428-35.</u></li> <li>Tamatani, T. <i>et al.</i> (2000) AlLIM/ICOS: a novel lymphocyte adhesion molecule. <u>Int Immunol. 12: 51-5.</u></li> <li>Dilek, N. <i>et al.</i> (2012) Control of transplant tolerance and intragraft regulatory T cell localization by myeloid-derived suppressor cells and CCL5. <u>J Immunol. 188: 4209-16.</u></li> <li>Ghiringhelli, F. <i>et al.</i> (2005) Tumor cells convert immature myeloid dendritic cells into TGF-beta-secreting cells inducing CD4+CD25+ regulatory T cell proliferation. <u>J Exp Med.</u> 202: 919-29.</li> <li>Sacedón, R. <i>et al.</i> (1999) Glucocorticoid-mediated regulation of thymic dendritic cell</li> </ol>			

	9. Kawai, T. <i>et al</i> .r (2000) T(h)1 transmigration anergy: a new concept of endothelial cell-T cell regulatory interaction. Int Immunol. 12: 937-48.				
	10. Macphee, I.A. et al. (2002) The Th2-response in mercuric chloride-induced				
	autoimmunity requires continuing costimulation via CD28. Clin Exp Immunol. 129: 405-10				
	11. MacPhee, I.A. <i>et al.</i> (2006) Blockade of OX40-ligand after initial triggering of the T				
	helper 2 response inhibits mercuric chloride-induced autoimmunity. Immunology. 117:				
	402-8.				
	12. Yrlid, U. <i>et al.</i> (2006) A distinct subset of intestinal dendritic cells responds selectively				
	to oral TLR7/8 stimulation. Eur J Immunol. 36: 2639-48.				
	13. Fan, C.B. et al. (2015) Alloantigen-specific T-cell hyporesponsiveness induced by				
	dnIKK2 gene-transfected recipient immature dendritic cells. Cell Immunol. 297 (2): 100-7.				
Storage	Store at +4°C or at -20°C if preferred.				
	Storage in frost-free freezers is not recommended.				
	This product should be stored undiluted. Avoid repeated freezing and thawing as this may				
	denature the antibody. Should this product contain a precipitate we recommend				
	microcentrifugation before use.				
Guarantee	18 months from date of despatch.				
Health And Safety	Material Safety Datasheet documentation #10040 available at:				
Information	10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf				
Regulatory	For research purposes only				

### **Related Products**

#### **Recommended Secondary Antibodies**

Goat Anti Mouse IgG IgA IgM (STAR87) <u>Alk. Phos.</u> , <u>HRP</u>			
Goat Anti Mouse IgG (STAR77)	HRP		
Rabbit Anti Mouse IgG (STAR12)	RPE		
Rabbit Anti Mouse IgG (STAR8)	DyLight®800		
Rabbit Anti Mouse IgG (STAR13)	HRP		
Goat Anti Mouse IgG (STAR76)	RPE		
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>		
Goat Anti Mouse IgG (Fc) (STAR120)	<u>FITC, HRP</u>		
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>		
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®680,		
	DyLight®800, FITC, HRP		

#### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL (MCA1209)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rac	l.com	Email: antibody_sales_uk@bio-rac	d.com	Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products.

#### Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint