Datasheet: MCA2876 BATCH NUMBER 154884

Description:	MOUSE ANTI HUMAN CA125		
Specificity:	CA125		
Other names:	MUCIN 16		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	X325		
Isotype:	lgG1		
Quantity:	0.2 mg		

Product Details

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry			•	
	Immunohistology - Frozen			•	
	Immunohistology - Paraffin	•			
	ELISA	•			
	Immunoprecipitation				
	Western Blotting	•			
	Where this antibody has	not been	tested fo	r use in a particular teo	hnique this does not
	necessarily exclude its us	se in suc	h procedu	res. Suggested workin	a dilutions are given as
	a guide only. It is recomm	ondod ti	hat tha us	or titratos the entibody	for use in their own
	a guide only. It is recomm	lended li	lat the us	er tittates the antibody	
	system using appropriate	negative	e/positive	controls.	
Target Species	Human				
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by	ion exch	ange chro	omatography	
Buffer Solution	TRIS buffered saline				

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)

Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	CA125 antigen purified from human ovarian carcinoma.
External Database Links	UniProt: Q8WXI7 Related reagents Entrez Gene: 94025 MUC16 Related reagents
Synonyms	CA125
RRID	AB_2075122
Fusion Partners	Spleen cells from immunised Balb/c mice were fused with cells of the Sp2/0 myeloma cell line.
Specificity	Mouse anti Human CA125 antibody, clone X325 recognizes the ovarian cancer mucin CA125 (also known as MUCIN 16). CA125 consists of a short cytoplasmic tail, a transmembrane domain and an exceptionally large glycosylated extracellular domain. The extracellular domain is dominated by a large number of 156-amino acid repeat units, over 60 in all. These repeats are known to bind the antibodies OC125 and M-11. CA125 is present within normal ovarian, endometrium, endocervix and fallopian tissue, but levels are elevated in over 90% of women with advanced ovarian cancer. CA125 levels may also be elevated in other cancers and non-cancerous conditions such as peritonitis, endometriosis and liver cirrhosis. Levels may also fluctuate during the menstrual cycle and occasionally increases in pregnancy, making CA125 an unreliable marker for ovarian cancer in premenopausal women. The function of CA125 in healthy tissue is unknown but it is thought to provide a protective lubricating barrier against particles and infectious agents at mucosal surfaces.
	CA125 was concluded by the International Society of Oncology and BioMarkers (ISOBM) to contain two major antigenic domains. Mouse anti Human CA125 antibody, clone X325 recognizes the epitope specificity group B, similar to the M-11 antibody (Nustad <i>et al.</i> 2002). Mouse anti Human CA125 antibody, clone X325 has been successfully as both a capture and detection reagent in the development of sensitive ELISA assays for the detection of CA125 antigen in human plasma samples.
Histology Positive Control Tissue	Lung adenocarcinoma, serosal ovarian adenocarcinoma and urinary bladder adenocarcinoma tissue.
References	1. Nustad, K. <i>et al.</i> (2002) Epitopes on CA 125 from cervical mucus and ascites fluid and characterization of six new antibodies. Third report from the ISOBM TD-1 workshop. <u>Tumour Biol. 23: 303-14.</u>
Storage	Store at +4°C or at -20°C if preferred.

	Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10057 available at: https://www.bio-rad-antibodies.com/SDS/MCA2876 10057
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	RPE
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>
Rabbit Anti Mouse IgG (STAR13)	HRP
Goat Anti Mouse IgG (Fc) (STAR120)	FITC, HRP
Rabbit Anti Mouse IgG (STAR9)	FITC
Goat Anti Mouse IgG (STAR77)	HRP
Goat Anti Mouse IgG (H/L) (STAR117)	<u>Alk. Phos.</u> , <u>DyLight®488</u> , <u>DyLight®550</u> ,
	DyLight®650, DyLight®680, DyLight®800,
	<u>FITC</u> , <u>HRP</u>

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@b	io-rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M367530:200529'

Printed on 14 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint