

Datasheet: MCA2877

Description:	Description: MOUSE ANTI HUMAN CA12	
Specificity:	CA125	
Other names:	MUCIN 16	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	X75	
Isotype:	lgG1	
Quantity:	0.2 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA	•			
Immunoprecipitation				
Western Blotting	•			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A	from ascites
Buffer Solution	150mM NaCl 50mM Citrate	
Preservative Stabilisers	≤0.1% Sodium Azide (NaN ₃)	

Approx. Protein Concentrations

IgG concentration 1.0mg/ml

Immunogen

CA125 antigen purified from human ovarian carcinoma.

External Database Links

UniProt:

Q8WXI7 Related reagents

Entrez Gene:

94025 MUC16 Related reagents

Synonyms

CA125

RRID

AB_2251124

Fusion Partners

Spleen cells from immunised Balb/c mice were fused with cells of the Sp2/0 myeloma cell line.

Specificity

Mouse anti Human CA125 antibody, clone X75 recognizes the ovarian cancer mucin CA125 (also known as MUCIN 16). CA125 consists of a short cytoplasmic tail, a transmembrane domain and an exceptionally large glycosylated extracellular domain. The extracellular domain is dominated by a large number of 156-amino acid repeat units, over 60 in all. These repeats are known to bind the antibodies OC125 and M-11.

CA125 is present within normal ovarian, endometrium, endocervix and fallopian tissue, but levels are elevated in over 90% of women with advanced ovarian cancer. CA125 levels may also be elevated in other cancers and non-cancerous conditions such as peritonitis, endometriosis and liver cirrhosis. Levels may also fluctuate during the menstrual cycle and occasionally increases in pregnancy, making CA125 an unreliable marker for ovarian cancer in premenopausal women. The function of CA125 in healthy tissue is unknown but it is thought to provide a protective lubricating barrier against particles and infectious agents at mucosal surfaces.

CA125 was concluded by the International Society of Oncology and BioMarkers (<u>ISOBM</u>) to contain two major antigenic domains. Mouse anti Human CA125 antibody, clone X75 recognizes the epitope specificity group B, similar to the M-11 antibody (<u>Nustad et al.</u> 2002). Mouse anti Human CA125 antibody, clone X75 has been used successfully as both a capture and detection reagent in the development of sensitive ELISA assays for the measurement of CA125 antigen in human plasma samples.

References

1. Nustad, K. *et al.* (2002) Epitopes on CA 125 from cervical mucus and ascites fluid and characterization of six new antibodies. Third report from the ISOBM TD-1 workshop. <u>Tumour Biol. 23: 303-14.</u>

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10196 available at: https://www.bio-rad-antibodies.com/SDS/MCA2877 10196
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...)

HRP

Goat Anti Mouse IgG (STAR70...)

Rabbit Anti Mouse IgG (STAR13...)

HRP

Rabbit Anti Mouse IgG (STAR9...)

Goat Anti Mouse IgG (STAR77...)

HRP

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376 Worl

America Fax: +1 919 878 3751

+1 800 265 7376 **Worldwide**

 ${\bf Email: antibody_sales_us@bio-rad.com}$

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M442823:250609'

Printed on 09 Jun 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint