

Datasheet: MCA2896PET

Description:	HAMSTER ANTI MOUSE CD178:RPE
Specificity:	CD178
Other names:	FAS LIGAND
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	MFL4
Isotype:	IgG
Quantity:	25 TEST

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	■			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Species Cross Reactivity	Reacts with: Rat N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized		
Reconstitution	Reconstitute with 0.25ml distilled water		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide (NaN ₃)		
Stabilisers	1% Bovine Serum Albumin		
	5% Sucrose		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml		

Immunogen	B6 mouse FasL/BHK cells
External Database Links	<p>UniProt: P41047 Related reagents</p> <p>Entrez Gene: 14103 FasL Related reagents</p>
Synonyms	Apt1lg1, Cd95l, Fasl, gld, Tnfsf6
Specificity	<p>Hamster anti Mouse CD178 antibody, clone MFL4 recognizes both mouse and rat CD178, otherwise known as Fas Ligand/ CD95L, a 40 kDa type II transmembrane glycoprotein and member of the TNF/NGF superfamily, expressed by activated T cells and NK cells, which can be induced on a variety of cells by radiation, heat shock, chemotherapeutic agents and viral infection.</p> <p>CD178 acts as a key effector of cytotoxicity and in the regulation of immune responses. The binding of CD178 to its receptor CD95 (Fas), induces Fas-mediated apoptosis of target cells, and may be involved in the induction of peripheral tolerance and neutrophil chemotaxis. The binding of decoy receptor 3 (DcR3) to CD178 has been shown to inhibit CD178-mediated apoptosis.</p> <p>Hamster anti Mouse CD178 antibody, clone MFL4 is reported to block CD178/CD95 induced apoptosis.</p>
Flow Cytometry	<p>Use 10ul of the suggested working dilution to label 1×10^6 cells in 100ul.</p> <p>The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B).</p>
References	<ol style="list-style-type: none"> 1. Trinité, B. <i>et al.</i> (2000) A subset of cytolytic dendritic cells in rat. J Immunol. 165 (8): 4202-8. 2. Watanabe, T. <i>et al.</i> (2002) Administration of an antigen at a high dose generates regulatory CD4+ T cells expressing CD95 ligand and secreting IL-4 in the liver. J Immunol. 168 (5): 2188-99. 3. Kayagaki, N. <i>et al.</i> (1997) Polymorphism of murine Fas ligand that affects the biological activity. Proc Natl Acad Sci U S A. 94 (8): 3914-9.
Storage	<p>Prior to reconstitution store at +4°C. After reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL:RPE \(MCA2356PE\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com
'M324302:180727'

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

Printed on 06 Dec 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)