

Datasheet: MCA329 BATCH NUMBER 158551

| Description: | MOUSE ANTI hCG (BETA 2 EPITOPE) | | |
|---------------|---------------------------------|--|--|
| Specificity: | hCG (BETA 2 EPITOPE) | | |
| Format: | Purified | | |
| Product Type: | Monoclonal Antibody | | |
| Clone: | INN-hCG-22 | | |
| Isotype: | IgG1 | | |
| Quantity: | 0.5 mg | | |
| | | | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|-------------------------|
| Flow Cytometry | | | • | |
| Immunohistology - Frozen | | | • | |
| Immunohistology - Paraffin | | | | |
| ELISA | | | | 1/100 - 1/500 |
| Immunoprecipitation | | | • | |
| Western Blotting | | | | Non-reducing conditions |
| Immunofluorescence | | | | |
| Immunoassay | | | | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| Target Species | Human |
|-----------------------------|---|
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide |
| Carrier Free | Yes |

| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml | | |
|--------------------------------|--|--|--|
| Immunogen | hCG. | | |
| External Database Links | UniProt: P01233 Related reagents | | |
| | Entrez Gene: 1082 CGB Related reagents | | |
| Synonyms | CGB3 | | |
| RRID | AB_2276154 | | |
| Fusion Partners | Spleen cells from an immunised BALB/c mouse were fused with cells of the mouse myeloma cell line P3 - NS1/1 - Ag 4.1. | | |
| Specificity | Mouse anti Human chorionic gonadotrophin antibody, clone INN-hCG-22 recognizes the beta subunit of human choriogonadotrophin (hCG), also known as chorionic gonadotrophin. hCGβ is a 165 amino acid ~18 kDa hormone involved in the stimulation of steroid production essential to the maintenance of pregnancy. | | |
| | Mouse anti Human chorionic gonadotrophin antibody, clone INN-hCG-22 shows a strong reaction in RIA with intact hCG and hCG β and some reactivity with human luteinizing hormone (12%) and b-hLH (34%). No reaction with human follicle-stimulating hormone, thyroid-stimulating hormone , a-hCG or a-hLH. | | |
| | Affinity constant = 1.6×10^9 m (Ka). | | |
| References | 1. Kofler, R. <i>et al.</i> (1982) Monoclonal antibodies against human chorionic gonadotropin (hCG): I. production, specificity, and intramolecular binding sites. <u>Am J Reprod Immunol. 2</u> (4): 212-6. | | |
| | 2. Schwarz, S. et al. (1985) Epitope-selective, monoclonal-antibody-based | | |

- immunoradiometric assays of predictable specificity for differential measurement of choriogonadotropin and its subunits. Clin Chem. 31 (8): 1322-8.
- 3. Zimmermann, G. et al. (2009) Epithelial human chorionic gonadotropin is expressed and produced in human secretory endometrium during the normal menstrual cycle. Biol Reprod. 80:1053-65.
- 4. Yoshie, M. et al. (2010) Possible role of the exchange protein directly activated by cyclic AMP (Epac) in the cyclic AMP-dependent functional differentiation and syncytialization of human placental BeWo cells. Hum Reprod. 25 (9): 2229-38.
- 5. Zimmermann, G. et al. (2003) Expression of beta hCG and alpha CG mRNA and hCG hormone in human decidual tissue in patients during tubal pregnancy. Mol Hum Reprod. 9: 81-9.
- 6. Birken, S. et al. (2003) Preparation and characterization of new WHO reference reagents for human chorionic gonadotropin and metabolites. Clin Chem. 49: 144-54.

- 7. Charrel-Dennis, M. *et al.* (2005) The human chorionic gonadotropin-beta arginine68 to glutamic acid substitution fixes the conformation of the C-terminal peptide. <u>Mol Endocrinol.</u> 19:1803-11.
- 8. Aye, I.L.M.H. *et al.* (2011) Oxysterols inhibit differentiation and fusion of term primary trophoblasts by activating liver X receptors <u>Placenta. 32: 183-91.</u>
- 9. Zimmermann, G. *et al.* (2012) Expression and production of human chorionic gonadotropin (hCG) in the normal secretory endometrium: evidence of CGB7 and/or CGB6 beta hCG subunit gene expression. Biol Reprod. 86 (3): 87.
- 10. Neuhaus, J. *et al.* (2011) New aspects in the differential diagnosis and therapy of bladder pain syndrome/interstitial cystitis. <u>Adv Urol.</u>;2011:639479.
- 11. Amantea, D. *et al.* (2016) Neuroprotective Properties of a Macrolide Antibiotic in a Mouse Model of Middle Cerebral Artery Occlusion: Characterization of the Immunomodulatory Effects and Validation of the Efficacy of Intravenous Administration. Assay Drug Dev Technol. 14 (5): 298-307.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

| Guarantee | 12 months from date of despatch |
|----------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA329 10040 |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...)

RPE

Goat Anti Mouse IgG (STAR76...)

RPE

Rabbit Anti Mouse IgG (STAR13...)

HRP

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_us@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M384039:210513'

Printed on 18 Jan 2024

@ 2024 Bio-Rad Laboratories Inc $\mid \underline{\text{Legal}} \mid \underline{\text{Imprint}}$