

Datasheet: MCA333S

BATCH NUMBER 1804

Description:	CHIMERIC HUMAN IgE ANTI NP
Specificity:	CHIMERIC HUMAN IgE ANTI NP
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	JW8/1
Isotype:	IgE
Quantity:	2 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/20 - 1/100
Immunoprecipitation			▪	
Western Blotting			▪	
Functional Assays (1)	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) This product contains sodium azide, removal by dialysis is recommended prior to use in functional assays.

Target Species	Human
Product Form	Tissue Culture Supernatant - liquid
Preparation	Tissue Culture Supernatant containing 0.2M Tris/HCl pH7.4 and 5-10% foetal calf serum
Buffer Solution	None present
Preservative Stabilisers	0.09% Sodium Azide

Immunogen	Hapten, 4-hydroxy-3-nitrophenylacetyl (NP).
RRID	AB_567284
Fusion Partners	Plasmids containing chimaeric heavy chain gene were fused with cells of the J558L mouse myeloma cell line.
Specificity	The immunoglobulin heavy chain has been produced by the linking of the antigen-binding, variable region genes of a mouse hybridoma to human constant region genes by <i>in vitro</i> DNA recombination procedures. The resulting chimeric antibody is subsequently expressed by the myeloma cell-line J558L after transfection. (The J558L cell-line self secretes a lambda light chain but no heavy chain). Thus a chimeric human IgE antibody specific for NP has been produced.
References	<ol style="list-style-type: none"> 1. Neuberger, M.S. <i>et al.</i> (1984) Recombinant antibodies possessing novel effector functions. Nature. 312 (5995): 604-8. 2. Sayers, I. <i>et al.</i> (2004) The importance of Lys-352 of human immunoglobulin E in FcepsilonRII/CD23 recognition. J Biol Chem. 279: 35320-5. 3. Neuberger, M.S. <i>et al.</i> (1985) A hapten-specific chimaeric IgE antibody with human physiological effector function. Nature. 314 (6008): 268-70. 4. Sallmann, E. <i>et al.</i> (2011) High-Affinity IgE Receptors on Dendritic Cells Exacerbate Th2-Dependent Inflammation. J Immunol. 187: 164-71. 5. Xu, D. <i>et al.</i> (2012) RN486, a selective Bruton's tyrosine kinase inhibitor, abrogates immune hypersensitivity responses and arthritis in rodents. J Pharmacol Exp Ther. 341 (1): 90-103. 6. Vangelista, L. <i>et al.</i> (2005) Membrane IgE binds and activates Fc epsilon RI in an antigen-independent manner. J Immunol. 174: 5602-11. 7. Kulka, M. and Metcalfe, D.D. (2004) High-resolution tracking of cell division demonstrates differential effects of TH1 and TH2 cytokines on SCF-dependent human mast cell production in vitro: correlation with apoptosis and Kit expression. Blood. 105: 592-9. 8. Novak, N. <i>et al.</i> (2003) Evidence for a differential expression of the FcepsilonRIgamma chain in dendritic cells of atopic and nonatopic donors. J Clin Invest. 111: 1047-56. 9. Sawada, J. <i>et al.</i> (2005) Stem cell factor has a suppressive activity to IgE-mediated chemotaxis of mast cells. J Immunol. 174: 3626-32. 10. Ferguson, G.D. <i>et al.</i> (2016) A Novel Triazolopyridine-Based Spleen Tyrosine Kinase Inhibitor That Arrests Joint Inflammation. PLoS One. 11 (1): e0145705. 11. Shirley D <i>et al.</i> (2016) Resveratrol preferentially inhibits IgE-dependent PGD₂ biosynthesis but enhances TNF production from human skin mast cells. Biochim Biophys Acta. pii: S0304-4165(16)00015-5. 12. Burton Oliver T. <i>et al.</i> (2016) A humanized mouse model of anaphylactic peanut allergy Journal of Allergy and Clinical Immunology. Jun 08 [Epub ahead of print] 13. Troupin, A. <i>et al.</i> (2016) A Role for Human Skin Mast Cells in Dengue Virus Infection and Systemic Spread. J Immunol. 197 (11): 4382-4391. 14. Bratke, K. <i>et al.</i> (2017) Differential regulation of PD-1 and its ligands in allergic asthma. Clin Exp Allergy. 47 (11): 1417-25. 15. Mchale, C.<i>et al.</i> (2018) Interleukin-6 potentiates FcεRI-induced PGD₂ biosynthesis and induces VEGF from human <i>in situ</i> -matured skin mast cells Biochimica et Biophysica

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10053 available at:
<https://www.bio-rad-antibodies.com/SDS/MCA333S>
10053

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M376477:210202'

Printed on 01 Aug 2024
