

Datasheet: MCA336

BATCH NUMBER 152980

Description:	RAT ANTI MOUSE IgG1 HEAVY CHAIN
Specificity:	IgG1 HEAVY CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	LO-MG1-2
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			5ug/ml as coating antibody
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide
Approx. Protein	IgG concentration 1 mg/ml

Concentrations

Immunogen Purified mouse IgG1 from BALB/c mice

External Database Links

UniProt:

[P01869](#) [Related reagents](#)

[P01868](#) [Related reagents](#)

Entrez Gene:

[16017](#) Ighg1 [Related reagents](#)

[16017](#) Ighg1 [Related reagents](#)

Synonyms Igh-4

RRID AB_321954

Fusion Partners Spleen cells from immunised LOU/c rats were fused with cells of the rat IR983F myeloma cell line.

Specificity **Rat anti Mouse IgG1 Heavy Chain antibody, clone LO-MG1-2** recognizes murine IgG1, and does not bind other mouse immunoglobulin classes or subclasses.

References

1. Song, J. *et al.* (2000) Heterogeneous distribution of isoactins in cultured vascular smooth muscle cells does not reflect segregation of contractile and cytoskeletal domains. [J Histochem Cytochem. 48 \(11\): 1441-52.](#)
2. Denis, O. *et al.* (1993) Resting B cells can act as antigen presenting cells in vivo and induce antibody responses. [Int Immunol. 5 \(1\): 71-8.](#)
3. Nakanishi, S. *et al.* (2010) Sequence analysis of a bacteriocinogenic plasmid of *Clostridium butyricum* and expression of the bacteriocin gene in *Escherichia coli*. [Anaerobe. 16: 253-7.](#)
4. Echeverria, P.C. *et al.* (2006) Potent antigen-specific immunity to *Toxoplasma gondii* in adjuvant-free vaccination system using Rop2-Leishmania infantum Hsp83 fusion protein. [Vaccine. 24: 4102-10.](#)
5. Huang, C.H. *et al.* (2011) Airway inflammation and IgE production induced by dust mite allergen-specific memory/effector Th2 cell line can be effectively attenuated by IL-35. [J Immunol. 187: 462-71.](#)
6. Agallou, M. *et al.* (2014) Experimental Validation of Multi-Epitope Peptides Including Promising MHC Class I- and II-Restricted Epitopes of Four Known *Leishmania infantum* Proteins. [Front Immunol. 5: 268.](#)
7. Doerfler, P.A. *et al.* (2015) BAFF Blockade Prevents Anti-Drug Antibody Formation in a Mouse Model of Pompe Disease. [Clin Immunol. pii: S1521-6616\(15\)00125-4.](#)
8. Ramos, J.D.A. *et al.* (2009) Characterization of Blo t 11 Monoclonal Antibodies with Constant Region Mutations *Phil Sci Lett.* 2(1): 38-48
9. Blackwell, N.M. & Else, K.J. (2002) A comparison of local and peripheral parasite-specific antibody production in different strains of mice infected with *Trichuris muris*. [Parasite Immunol. 24 \(4\): 203-11.](#)
10. Hall, G. *et al.* (2003) Suppression of allergen reactive Th2 mediated responses and

pulmonary eosinophilia by intranasal administration of an immunodominant peptide is linked to IL-10 production. [Vaccine. 21 \(5-6\): 549-61.](#)

11. Hjerpe, C. *et al.* (2010) Dendritic cells pulsed with malondialdehyde modified low density lipoprotein aggravate atherosclerosis in Apoe(-/-) mice. [Atherosclerosis. 209 \(2\): 436-41.](#)

12. Kretschmer, B. *et al.* (2015) Anti-CD83 promotes IgG1 isotype switch in marginal zone B cells in response to TI-2 antigen. [Immunobiology. 220 \(8\): 964-75.](#)

13. Doerfler, P.A. *et al.* (2016) Copackaged AAV9 Vectors Promote Simultaneous Immune Tolerance and Phenotypic Correction of Pompe Disease. [Hum Gene Ther. 27 \(1\): 43-59.](#)

14. Kato, G. *et al.* (2014) β 2 adrenergic agonist attenuates house dust mite-induced allergic airway inflammation through dendritic cells. [BMC Immunol. 15: 39.](#)

15. Margaroni, M. *et al.* (2017) Vaccination with poly(D,L-lactide-co-glycolide) nanoparticles loaded with soluble *Leishmania* antigens and modified with a TNF α -mimicking peptide or monophosphoryl lipid A confers protection against experimental visceral leishmaniasis. [Int J Nanomedicine. 12: 6169-84.](#)

16. DeGiovanni, C. *et al.* (2019) Cancer Vaccines Co-Targeting HER2/Neu and IGF1R. [Cancers \(Basel\). 11 \(4\) Apr 11 \[Epub ahead of print\].](#)

17. Doshi, B.S. *et al.* (2021) B cell-activating factor modulates the factor VIII immune response in hemophilia A. [J Clin Invest. 131\(8\):142906.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA336>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

North & South Tel: +1 800 265 7376

Worldwide Tel: +44 (0)1865 852 700

Europe Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M367584:200529'

Printed on 25 Mar 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)