

Datasheet: MCA336

Description:	RAT ANTI MOUSE IgG1 HEAVY CHAIN
Specificity:	IgG1 HEAVY CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	LO-MG1-2
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

RRID AB_321954

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			5ug/ml as coating antibody
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Mouse

Product Form Purified IgG - liquid

Preparation Purified IgG prepared by affinity chromatography from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.1% Sodium Azide

Approx. Protein Concentrations IgG concentration 1 mg/ml

Immunogen Purified mouse IgG1 from BALB/c mice

External Database **UniProt:**

Links[P01869](#) [Related reagents](#)[P01868](#) [Related reagents](#)**Entrez Gene:**[16017](#) Ighg1 [Related reagents](#)[16017](#) Ighg1 [Related reagents](#)**Synonyms**

Igh-4

Fusion Partners

Spleen cells from immunised LOU/c rats were fused with cells of the rat IR983F myeloma cell line.

Specificity**Rat anti Mouse IgG1 Heavy Chain antibody, clone LO-MG1-2** recognizes murine IgG1, and does not bind other mouse immunoglobulin classes or subclasses.**References**

1. Song, J. *et al.* (2000) Heterogeneous distribution of isoactins in cultured vascular smooth muscle cells does not reflect segregation of contractile and cytoskeletal domains. [J Histochem Cytochem. 48 \(11\): 1441-52.](#)
2. Denis, O. *et al.* (1993) Resting B cells can act as antigen presenting cells in vivo and induce antibody responses. [Int Immunol. 5 \(1\): 71-8.](#)
3. Nakanishi, S. *et al.* (2010) Sequence analysis of a bacteriocinogenic plasmid of *Clostridium butyricum* and expression of the bacteriocin gene in *Escherichia coli*. [Anaerobe. 16: 253-7.](#)
4. Echeverria, P.C. *et al.* (2006) Potent antigen-specific immunity to *Toxoplasma gondii* in adjuvant-free vaccination system using Rop2-Leishmania infantum Hsp83 fusion protein. [Vaccine. 24: 4102-10.](#)
5. Huang, C.H. *et al.* (2011) Airway inflammation and IgE production induced by dust mite allergen-specific memory/effector Th2 cell line can be effectively attenuated by IL-35. [J Immunol. 187: 462-71.](#)
6. Agallou, M. *et al.* (2014) Experimental Validation of Multi-Epitope Peptides Including Promising MHC Class I- and II-Restricted Epitopes of Four Known *Leishmania infantum* Proteins. [Front Immunol. 5: 268.](#)
7. Doerfler, P.A. *et al.* (2015) BAFF Blockade Prevents Anti-Drug Antibody Formation in a Mouse Model of Pompe Disease. [Clin Immunol. pii: S1521-6616\(15\)00125-4.](#)
8. Ramos, J.D.A. *et al.* (2009) Characterization of Blo t 11 Monoclonal Antibodies with Constant Region Mutations [Phil Sci Lett. 2\(1\): 38-48.](#)
9. Blackwell, N.M. & Else, K.J. (2002) A comparison of local and peripheral parasite-specific antibody production in different strains of mice infected with *Trichuris muris*. [Parasite Immunol. 24 \(4\): 203-11.](#)
10. Hall, G. *et al.* (2003) Suppression of allergen reactive Th2 mediated responses and pulmonary eosinophilia by intranasal administration of an immunodominant peptide is linked to IL-10 production. [Vaccine. 21 \(5-6\): 549-61.](#)
11. Hjerpe, C. *et al.* (2010) Dendritic cells pulsed with malondialdehyde modified low density lipoprotein aggravate atherosclerosis in Apoe(-/-) mice. [Atherosclerosis. 209 \(2\): 436-41.](#)
12. Kretschmer, B. *et al.* (2015) Anti-CD83 promotes IgG1 isotype switch in marginal zone B cells in response to TI-2 antigen. [Immunobiology. 220 \(8\): 964-75.](#)
13. Doerfler, P.A. *et al.* (2016) Copackaged AAV9 Vectors Promote Simultaneous Immune Tolerance and Phenotypic Correction of Pompe Disease. [Hum Gene Ther. 27 \(1\): 43-59.](#)
14. Kato, G. *et al.* (2014) β 2 adrenergic agonist attenuates house dust mite-induced allergic airway inflammation through dendritic cells. [BMC Immunol. 15: 39.](#)
15. Margaroni, M. *et al.* (2017) Vaccination with poly(D,L-lactide-co-glycolide) nanoparticles loaded with soluble *Leishmania* antigens and modified with a TNF α -mimicking peptide or monophosphoryl lipid A confers protection against experimental visceral leishmaniasis. [Int J Nanomedicine. 12:](#)

[6169-84.](#)

16. DeGiovanni, C. *et al.* (2019) Cancer Vaccines Co-Targeting HER2/Neu and IGF1R. [Cancers \(Basel\). 11 \(4\) Apr 11 \[Epub ahead of print\].](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10303 available at:
10303: <https://www.bio-rad-antibodies.com/uploads/MSDS/10303.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (STAR72...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®649 , DyLight®800

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M339525:181219'

Printed on 16 Apr 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)