

Datasheet: MCA336

Description:	RAT ANTI MOUSE IgG1 HEAVY CHAIN
Specificity:	IgG1 HEAVY CHAIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	LO-MG1-2
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			5ug/ml as coating antibody
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1 mg/ml

Immunogen Purified mouse IgG1 from BALB/c mice

External Database

Links

UniProt:

[P01869](#) [Related reagents](#)

[P01868](#) [Related reagents](#)

Entrez Gene:

[16017](#) Ighg1 [Related reagents](#)

[16017](#) Ighg1 [Related reagents](#)

Synonyms Igh-4

RRID AB_321954

Fusion Partners Spleen cells from immunised LOU/c rats were fused with cells of the rat IR983F myeloma cell line.

Specificity **Rat anti Mouse IgG1 Heavy Chain antibody, clone LO-MG1-2** recognizes murine IgG1, and does not bind other mouse immunoglobulin classes or subclasses.

References

1. Song, J. *et al.* (2000) Heterogeneous distribution of isoactins in cultured vascular smooth muscle cells does not reflect segregation of contractile and cytoskeletal domains. [J Histochem Cytochem. 48 \(11\): 1441-52.](#)
2. Denis, O. *et al.* (1993) Resting B cells can act as antigen presenting cells in vivo and induce antibody responses. [Int Immunol. 5 \(1\): 71-8.](#)
3. Nakanishi, S. *et al.* (2010) Sequence analysis of a bacteriocinogenic plasmid of *Clostridium butyricum* and expression of the bacteriocin gene in *Escherichia coli*. [Anaerobe. 16: 253-7.](#)
4. Echeverria, P.C. *et al.* (2006) Potent antigen-specific immunity to *Toxoplasma gondii* in adjuvant-free vaccination system using Rop2-Leishmania infantum Hsp83 fusion protein. [Vaccine. 24: 4102-10.](#)
5. Huang, C.H. *et al.* (2011) Airway inflammation and IgE production induced by dust mite allergen-specific memory/effector Th2 cell line can be effectively attenuated by IL-35. [J Immunol. 187: 462-71.](#)
6. Agallou, M. *et al.* (2014) Experimental Validation of Multi-Epitope Peptides Including Promising MHC Class I- and II-Restricted Epitopes of Four Known *Leishmania infantum* Proteins. [Front Immunol. 5: 268.](#)
7. Doerfler, P.A. *et al.* (2015) BAFF Blockade Prevents Anti-Drug Antibody Formation in a Mouse Model of Pompe Disease. [Clin Immunol. pii: S1521-6616\(15\)00125-4.](#)
8. Ramos, J.D.A. *et al.* (2009) Characterization of 11 Monoclonal Antibodies with Constant Region Mutations *Phil Sci Lett.* 2(1): 38-48
9. Blackwell, N.M. & Else, K.J. (2002) A comparison of local and peripheral parasite-specific antibody production in different strains of mice infected with *Trichuris muris*. [Parasite Immunol. 24 \(4\): 203-11.](#)
10. Hall, G. *et al.* (2003) Suppression of allergen reactive Th2 mediated responses and pulmonary eosinophilia by intranasal administration of an immunodominant peptide is linked to IL-10 production. [Vaccine. 21 \(5-6\): 549-61.](#)

11. Hjerpe, C. *et al.* (2010) Dendritic cells pulsed with malondialdehyde modified low density lipoprotein aggravate atherosclerosis in Apoe(-/-) mice. [Atherosclerosis. 209 \(2\): 436-41.](#)
12. Kretschmer, B. *et al.* (2015) Anti-CD83 promotes IgG1 isotype switch in marginal zone B cells in response to TI-2 antigen. [Immunobiology. 220 \(8\): 964-75.](#)
13. Doerfler, P.A. *et al.* (2016) Copackaged AAV9 Vectors Promote Simultaneous Immune Tolerance and Phenotypic Correction of Pompe Disease. [Hum Gene Ther. 27 \(1\): 43-59.](#)
14. Kato, G. *et al.* (2014) β 2 adrenergic agonist attenuates house dust mite-induced allergic airway inflammation through dendritic cells. [BMC Immunol. 15: 39.](#)
15. Margaroni, M. *et al.* (2017) Vaccination with poly(D,L-lactide-co-glycolide) nanoparticles loaded with soluble *Leishmania* antigens and modified with a TNF α -mimicking peptide or monophosphoryl lipid A confers protection against experimental visceral leishmaniasis. [Int J Nanomedicine. 12: 6169-84.](#)
16. DeGiovanni, C. *et al.* (2019) Cancer Vaccines Co-Targeting HER2/Neu and IGF1R. [Cancers \(Basel\). 11 \(4\) Apr 11 \[Epub ahead of print\].](#)
17. Doshi, B.S. *et al.* (2021) B cell-activating factor modulates the factor VIII immune response in hemophilia A. [J Clin Invest. 131\(8\):142906.](#)
18. Scaramozza, A. *et al.* (2019) Lineage Tracing Reveals a Subset of Reserve Muscle Stem Cells Capable of Clonal Expansion under Stress. [Cell Stem Cell. 24 \(6\): 944-957.e5.](#)
19. Steel, N. *et al.* (2019) TGF β -activation by dendritic cells drives Th17 induction and intestinal contractility and augments the expulsion of the parasite *Trichinella spiralis* in mice. [PLoS Pathog. 15 \(4\): e1007657.](#)
20. Salem, E. *et al.* (2019) Pathogenesis, Host Innate Immune Response, and Aerosol Transmission of Influenza D Virus in Cattle. [J Virol.93\(7\):e01853-18.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR16...)	DyLight@800
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@650 , DyLight@800
Goat Anti Rat IgG (STAR72...)	HRP

Rabbit Anti Rat IgG (STAR21...)

[HRP](#)

Rabbit Anti Rat IgG (STAR17...)

[FITC](#)

Goat Anti Rat IgG (STAR131...)

[Alk. Phos.](#), [Biotin](#)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M384492:210513'

Printed on 21 Mar 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)