

## Datasheet: MCA336B

**BATCH NUMBER 167056**

<b>Description:</b>	RAT ANTI MOUSE IgG1 HEAVY CHAIN:Biotin
<b>Specificity:</b>	IgG1 HEAVY CHAIN
<b>Format:</b>	Biotin
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	LO-MG1-2
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.5 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			5 ug/ml
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG conjugated to Biotin - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.1% Sodium Azide 50% Glycerol
<b>Approx. Protein Concentrations</b>	IgG concentration 1 mg/ml

Immunogen Purified mouse IgG1 from BALB/c mice

---

**External Database**

**Links**

**UniProt:**

[P01869](#) [Related reagents](#)

[P01868](#) [Related reagents](#)

**Entrez Gene:**

[16017](#) Ighg1 [Related reagents](#)

[16017](#) Ighg1 [Related reagents](#)

---

**Synonyms** Igh-4

---

**RRID** AB\_321955

---

**Fusion Partners** Spleen cells from immunised LOU/c rats were fused with cells of the rat IR983F myeloma cell line.

---

**Specificity** **Rat anti Mouse IgG1 Heavy Chain antibody, clone LO-MG1-2** recognizes murine IgG1, and does not bind other mouse immunoglobulin classes or subclasses.

---

**References**

1. Denis, O. *et al.* (1993) Resting B cells can act as antigen presenting cells in vivo and induce antibody responses. [Int Immunol. 5 \(1\): 71-8.](#)
2. Song, J. *et al.* (2000) Heterogeneous distribution of isoactins in cultured vascular smooth muscle cells does not reflect segregation of contractile and cytoskeletal domains. [J Histochem Cytochem. 48 \(11\): 1441-52.](#)
3. Blackwell, N.M. & Else, K.J. (2002) A comparison of local and peripheral parasite-specific antibody production in different strains of mice infected with *Trichuris muris*. [Parasite Immunol. 24 \(4\): 203-11.](#)
4. Hall, G. *et al.* (2003) Suppression of allergen reactive Th2 mediated responses and pulmonary eosinophilia by intranasal administration of an immunodominant peptide is linked to IL-10 production. [Vaccine. 21 \(5-6\): 549-61.](#)
5. Echeverria, P.C. *et al.* (2006) Potent antigen-specific immunity to *Toxoplasma gondii* in adjuvant-free vaccination system using Rop2-*Leishmania infantum* Hsp83 fusion protein. [Vaccine. 24: 4102-10.](#)
6. Ramos, J.D.A. *et al.* (2009) Characterization of Blo t 11 Monoclonal Antibodies with Constant Region Mutations Phil Sci Lett. 2(1): 38-48
7. Nakanishi, S. *et al.* (2010) Sequence analysis of a bacteriocinogenic plasmid of *Clostridium butyricum* and expression of the bacteriocin gene in *Escherichia coli*. [Anaerobe. 16: 253-7.](#)
8. Hjerpe, C. *et al.* (2010) Dendritic cells pulsed with malondialdehyde modified low density lipoprotein aggravate atherosclerosis in Apoe(-/-) mice. [Atherosclerosis. 209 \(2\): 436-41.](#)
9. Huang, C.H. *et al.* (2011) Airway inflammation and IgE production induced by dust mite allergen-specific memory/effector Th2 cell line can be effectively attenuated by IL-35. [J Immunol. 187: 462-71.](#)
10. Agallou, M. *et al.* (2014) Experimental Validation of Multi-Epitope Peptides Including Promising MHC Class I- and II-Restricted Epitopes of Four Known *Leishmania infantum*

Proteins. [Front Immunol. 5: 268.](#)

11. Kato, G. *et al.* (2014)  $\beta$ 2 adrenergic agonist attenuates house dust mite-induced allergic airway inflammation through dendritic cells. [BMC Immunol. 15: 39.](#)
12. Doerfler, P.A. *et al.* (2015) BAFF Blockade Prevents Anti-Drug Antibody Formation in a Mouse Model of Pompe Disease. [Clin Immunol. pii: S1521-6616\(15\)00125-4.](#)
13. Kretschmer, B. *et al.* (2015) Anti-CD83 promotes IgG1 isotype switch in marginal zone B cells in response to TI-2 antigen. [Immunobiology. 220 \(8\): 964-75.](#)
14. Doerfler, P.A. *et al.* (2016) Copackaged AAV9 Vectors Promote Simultaneous Immune Tolerance and Phenotypic Correction of Pompe Disease. [Hum Gene Ther. 27 \(1\): 43-59.](#)
15. Margaroni, M. *et al.* (2017) Vaccination with poly(D,L-lactide-co-glycolide) nanoparticles loaded with soluble *Leishmania* antigens and modified with a TNF $\alpha$ -mimicking peptide or monophosphoryl lipid A confers protection against experimental visceral leishmaniasis. [Int J Nanomedicine. 12: 6169-84.](#)
16. Steel, N. *et al.* (2019) TGF $\beta$ -activation by dendritic cells drives Th17 induction and intestinal contractility and augments the expulsion of the parasite *Trichinella spiralis* in mice. [PLoS Pathog. 15 \(4\): e1007657.](#)
17. DeGiovanni, C. *et al.* (2019) Cancer Vaccines Co-Targeting HER2/Neu and IGF1R. [Cancers \(Basel\). 11 \(4\) Apr 11 \[Epub ahead of print\].](#)
18. Scaramozza, A. *et al.* (2019) Lineage Tracing Reveals a Subset of Reserve Muscle Stem Cells Capable of Clonal Expansion under Stress. [Cell Stem Cell. 24 \(6\): 944-957.e5.](#)
19. Salem, E. *et al.* (2019) Pathogenesis, Host Innate Immune Response, and Aerosol Transmission of Influenza D Virus in Cattle. [J Virol. 93\(7\):e01853-18.](#)
20. Doshi, B.S. *et al.* (2021) B cell-activating factor modulates the factor VIII immune response in hemophilia A. [J Clin Invest. 131\(8\):142906.](#)
21. Petta, I. *et al.* (2024) Myeloid A20 is critical for alternative macrophage polarization and type-2 immune mediated helminth resistance [Front Immunol. 15: 21 Mar \[Epub ahead of print\]](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10328 available at: <https://www.bio-rad-antibodies.com/SDS/MCA336B>  
10328

---

**Regulatory**

For research purposes only

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M384040:210513'

