

Datasheet: MCA336P

Description:	RAT ANTI MOUSE IgG1 HEAVY CHAIN:HRP
Specificity:	IgG1 HEAVY CHAIN
Format:	HRP
Product Type:	Monoclonal Antibody
Clone:	LO-MG1-2
Isotype:	IgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			500ng/ml
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG conjugated to Horseradish Peroxidase (HRP) - liquid
Preparation	Purified IgG prepared by affinity chromatography from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.01% Thiomersal 50% Glycerol
Approx. Protein Concentrations	IgG concentration 1 mg/ml
Immunogen	Purified mouse IgG1 from BALB/c mice

External Database
Links

UniProt:

[P01869](#) [Related reagents](#)
[P01868](#) [Related reagents](#)

Entrez Gene:

[16017](#) Ighg1 [Related reagents](#)
[16017](#) Ighg1 [Related reagents](#)

Synonyms	Igh-4
-----------------	-------

RRID	AB_321957
-------------	-----------

Fusion Partners	Spleen cells from immunised LOU/c rats were fused with cells of the rat IR983F myeloma cell line.
------------------------	---

Specificity	Rat anti Mouse IgG1 Heavy Chain antibody, clone LO-MG1-2 recognizes murine IgG1, and does not bind other mouse immunoglobulin classes or subclasses.
--------------------	---

ELISA	This product may be used in a direct ELISA or as a detection reagent in a sandwich ELISA together with MCA1289 as the capture reagent.
--------------	--

References	<ol style="list-style-type: none">1. Song, J. <i>et al.</i> (2000) Heterogeneous distribution of isoactins in cultured vascular smooth muscle cells does not reflect segregation of contractile and cytoskeletal domains. J Histochem Cytochem. 48 (11): 1441-52.2. Denis, O. <i>et al.</i> (1993) Resting B cells can act as antigen presenting cells in vivo and induce antibody responses. Int Immunol. 5 (1): 71-8.3. Nakanishi, S. <i>et al.</i> (2010) Sequence analysis of a bacteriocinogenic plasmid of <i>Clostridium butyricum</i> and expression of the bacteriocin gene in <i>Escherichia coli</i>. Anaerobe. 16: 253-7.4. Echeverria, P.C. <i>et al.</i> (2006) Potent antigen-specific immunity to <i>Toxoplasma gondii</i> in adjuvant-free vaccination system using Rop2-Leishmania infantum Hsp83 fusion protein. Vaccine. 24: 4102-10.5. Huang, C.H. <i>et al.</i> (2011) Airway inflammation and IgE production induced by dust mite allergen-specific memory/effector Th2 cell line can be effectively attenuated by IL-35. J Immunol. 187: 462-71.6. Agallou, M. <i>et al.</i> (2014) Experimental Validation of Multi-Epitope Peptides Including Promising MHC Class I- and II-Restricted Epitopes of Four Known <i>Leishmania infantum</i> Proteins. Front Immunol. 5: 268.7. Doerfler, P.A. <i>et al.</i> (2015) BAFF Blockade Prevents Anti-Drug Antibody Formation in a Mouse Model of Pompe Disease. Clin Immunol. pii: S1521-6616(15)00125-4.8. Ramos, J.D.A. <i>et al.</i> (2009) Characterization of Blo t 11 Monoclonal Antibodies with Constant Region Mutations Phil Sci Lett. 2(1): 38-489. Blackwell, N.M. & Else, K.J. (2002) A comparison of local and peripheral parasite-specific antibody production in different strains of mice infected with <i>Trichuris muris</i>. Parasite Immunol. 24 (4): 203-11.10. Hall, G. <i>et al.</i> (2003) Suppression of allergen reactive Th2 mediated responses and pulmonary eosinophilia by intranasal administration of an immunodominant peptide is
-------------------	--

linked to IL-10 production. [Vaccine. 21 \(5-6\): 549-61.](#)

11. Hjerpe, C. *et al.* (2010) Dendritic cells pulsed with malondialdehyde modified low density lipoprotein aggravate atherosclerosis in Apoe(-/-) mice. [Atherosclerosis. 209 \(2\): 436-41.](#)

12. Kretschmer, B. *et al.* (2015) Anti-CD83 promotes IgG1 isotype switch in marginal zone B cells in response to TI-2 antigen. [Immunobiology. 220 \(8\): 964-75.](#)

13. Doerfler, P.A. *et al.* (2016) Copackaged AAV9 Vectors Promote Simultaneous Immune Tolerance and Phenotypic Correction of Pompe Disease. [Hum Gene Ther. 27 \(1\): 43-59.](#)

14. Kato, G. *et al.* (2014) β 2 adrenergic agonist attenuates house dust mite-induced allergic airway inflammation through dendritic cells. [BMC Immunol. 15: 39.](#)

15. Margaroni, M. *et al.* (2017) Vaccination with poly(D,L-lactide-co-glycolide) nanoparticles loaded with soluble *Leishmania* antigens and modified with a TNF α -mimicking peptide or monophosphoryl lipid A confers protection against experimental visceral leishmaniasis. [Int J Nanomedicine. 12: 6169-84.](#)

16. DeGiovanni, C. *et al.* (2019) Cancer Vaccines Co-Targeting HER2/Neu and IGF1R. [Cancers \(Basel\). 11 \(4\) Apr 11 \[Epub ahead of print\].](#)

17. Doshi, B.S. *et al.* (2021) B cell-activating factor modulates the factor VIII immune response in hemophilia A. [J Clin Invest. 131\(8\):142906.](#)

18. Scaramozza, A. *et al.* (2019) Lineage Tracing Reveals a Subset of Reserve Muscle Stem Cells Capable of Clonal Expansion under Stress. [Cell Stem Cell. 24 \(6\): 944-957.e5.](#)

19. Steel, N. *et al.* (2019) TGF β -activation by dendritic cells drives Th17 induction and intestinal contractility and augments the expulsion of the parasite *Trichinella spiralis* in mice. [PLoS Pathog. 15 \(4\): e1007657.](#)

20. Salem, E. *et al.* (2019) Pathogenesis, Host Innate Immune Response, and Aerosol Transmission of Influenza D Virus in Cattle. [J Virol. 93\(7\):e01853-18.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10097 available at: 10097: <https://www.bio-rad-antibodies.com/uploads/MSDS/10097.pdf>

Regulatory

For research purposes only

Related Products

Recommended Useful Reagents

[AbGUARD® HRP STABILIZER PLUS \(BUF052A\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052B\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052C\)](#)

[TMB CORE \(BUF056A\)](#)

[TMB CORE+ \(BUF062A\)](#)

[TMB SIGNAL+ \(BUF054A\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M384066:210513'

Printed on 21 Mar 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)